

bs-2388R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

GPD1 Rabbit pAb**DATASHEET**

| | | |
|--|----------------------|--|
| Host: Rabbit | Isotype: IgG | Applications: WB (1:500-2000) |
| Clonality: Polyclonal | | |
| GeneID: 2819 | SWISS: P21695 | |
| Target: GPD1 | | |
| Immunogen: KLH conjugated synthetic peptide derived from human GPD1: 51-150/349. | | |
| Purification: affinity purified by Protein A | | Reactivity: Mouse, Rat (predicted: Human, Rabbit, Pig, Cow, Horse) |
| Concentration: 1mg/ml | | |
| Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. | | |
| Background: Glycerol 3 Phosphate Dehydrogenase is a homodimer and belongs to the NAD dependent Glycerol 3 Phosphate Dehydrogenase family. | | |
| | | |
| | | Predicted MW.: 37 kDa |
| | | Subcellular Location: Cytoplasm |

VALIDATION IMAGES

Sample: Lane 1: Mouse Liver tissue lysates Lane 2: Rat Liver tissue lysates Primary: Anti-GPD1 (bs-2388R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 37 kDa Observed band size: 37 kDa

SELECTED CITATIONS

- **[IF=13.6]** Shuping Zhong. et al. Persistent metabolic toxicities following developmental exposure to hexafluoropropylene oxide trimer acid (HFPO-TA): Roles of peroxisome proliferator activated receptor gamma. J HAZARD MATER. 2024 Jun;471:134337 IHC ;Chicken. 38640674
- **[IF=4.379]** Li, Shen. et al. Transcriptome profile of goat folliculogenesis reveals the interaction of oocyte and granulosa cell in correlation with different fertility population. Sci Rep-Uk. 2021 Aug;11(1):1-14 IF,IHC ;Goat. 34344973