

**bs-11689R****[ Primary Antibody ]****FGGY Rabbit pAb****BioSS**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**— DATASHEET —**

<b>Host:</b> Rabbit <b>Clonality:</b> Polyclonal <b>GeneID:</b> 55277 <b>Target:</b> FGGY <b>Immunogen:</b> KLH conjugated synthetic peptide derived from human FGGY: 151-250/551. <b>Purification:</b> affinity purified by Protein A <b>Concentration:</b> 1mg/ml <b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. <b>Background:</b> FGGY is a 551 amino acid member of the FGGY kinase family that exists as four isoforms which are produced by alternative splicing events. Expressed in lung, kidney, small intestine, liver and fetal brain, FGGY is encoded by a gene that maps to chromosome 1 and, when mutated, is associated with sporadic amyotrophic lateral sclerosis (ALS). ALS is a neurodegenerative disorder that affects motor neurons and results in fatal paralysis, usually within 2 to 5 years after initial diagnosis. Chromosome 1, on which the gene encoding FGGY is located, is the largest human chromosome, spanning about 260 million base pairs and making up 8% of the human genome. There are about 3,000 genes on chromosome 1, many of which are associated with genetic diseases, including Hutchinson-Gilford progeria, familial adenomatous polyposis, Stickler syndrome, Gaucher disease and Usher syndrome.	<b>Isotype:</b> IgG <b>SWISS:</b> Q96C11 <b>Applications:</b> IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) <b>Reactivity:</b> Mouse, Rat (predicted: Human, Rabbit, Sheep, Cow, Dog, Horse) <b>Predicted MW.:</b> 60 kDa <b>Subcellular Location:</b> Cytoplasm ,Nucleus
--	--

**— VALIDATION IMAGES —**

Paraformaldehyde-fixed, paraffin embedded (mouse liver tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FGGY) Polyclonal Antibody, Unconjugated (bs-11689R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat kidney tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FGGY) Polyclonal Antibody, Unconjugated (bs-11689R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.