

**bs-18436R****[ Primary Antibody ]****LSM12 Rabbit pAb****BioSS**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**— DATASHEET —**

|  |                      |   |
|--|----------------------|---|
| <b>Host:</b> Rabbit  | <b>Isotype:</b> IgG  | <b>Applications:</b> IHC-P (1:100-500)<br>IHC-F (1:100-500)<br>IF (1:100-500)<br><br><b>Reactivity:</b> Mouse, Rat<br>(predicted: Human, Rabbit, Pig)<br><br><b>Predicted MW.:</b> 22 kDa<br><br><b>Subcellular Location:</b> Nucleus |
| <b>Clonality:</b> Polyclonal   |                      |   |
| <b>GeneID:</b> 124801  | <b>SWISS:</b> Q3MHD2 |   |
| <b>Target:</b> LSM12   |                      |   |
| <b>Immunogen:</b> KLH conjugated synthetic peptide derived from human LSM12: 101-195/195.  |                      |   |
| <b>Purification:</b> affinity purified by Protein A  |                      |   |
| <b>Concentration:</b> 1mg/ml   |                      |   |
| <b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.<br>Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. |                      |   |
| <b>Background:</b> LSM12 belongs to the LSM12 family. The exact function of LSM12 is not known.  |                      |   |

**— VALIDATION IMAGES —**

Paraformaldehyde-fixed, paraffin embedded (rat liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (LSM12) Polyclonal Antibody, Unconjugated (bs-18436R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (LSM12) Polyclonal Antibody, Unconjugated (bs-18436R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.