bs-9001R

[Primary Antibody]

SARM1 Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET		400-901-9800
Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500)
Clonality: Polyclonal		IHC-F (1:100-500) IF (1:50-200)
GenelD: 23098	SWISS: Q6SZW1	
Target: SARM1		Reactivity: Human (predicted: Mouse, Rat Rabbit Pig Cow Dog
Immunogen: KLH conjugated synthetic peptide derived from human SARM1: 321-420/724.		Horse)
Purification: affinity purified by Protein A		Predicted 80 kDa
Concentration: 1mg/ml		MW.: OUNDU
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Location: ^{Cytoplasm}
Background: SARM, also known as SAMD2, SARM1 or KIAA0524, is a 724 amino acid protein that localizes to the cytoplasm and contains one TIR domain and two sterile alpha motif (SAM) domains. Expressed predominately in liver and kidney and present at lower levels in placenta, SARM interacts with TICAM-1 and, via this interaction, blocks the transcriptional activation activity of TICAM-1 and functions as a negative regulator of Toll-like receptor signaling. Additionally, SARM is thought to be involved in innate immune responses and may also play a role in the negative regulation of NF x B activation. SARM exists as two alternatively spliced isoforms that are encoded by a gene which maps to human chromosome 17.		

- VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (Human liver tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (SARM1) Polyclonal Antibody, Unconjugated (bs-9001R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.

- SELECTED CITATIONS -

• [IF=2.992] Cuiqin Zhang. et al. Atg7 Knockout Alleviated the Axonal Injury of Neuro-2a Cells Induced by Tri-Ortho-Cresyl Phosphate. 2021 Mar 01 WB ;Mouse. 33650059