### bs-3507R

- DATASHEET -

## [ Primary Antibody ]

# NMDAR2A Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHE	- '		
Host	Rabbit	Isotype: IgG	Applications: WB (1:500-1:2000)
Clonality: Polyclonal			<b>Reactivity:</b> Human (predicted: Mouse.
GenelD:	2903	SWISS: Q12879	Rat)
Target:	NMDAR2A		
Immunogen: KLH conjugated synthetic peptide derived from human NMDAR2A: 851-950/1462.			Predicted MW.: <sup>163 kDa</sup>
Purification: affinity purified by Protein A			Cubacillular
Concentration: 1mg/ml			Location: Cell membrane
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.			
<b>Background:</b> N-methyl-D-aspartate (NMDA) receptors are a class of ionotropic glutamate-gated ion channels. These receptors have been shown to be involved in long-term potentiation, an activity-dependent increase in the efficiency of synaptic transmission thought to underlie certain kinds of memory and learning. NMDA receptor channels are heteromers composed of the key receptor subunit NMDAR1 (GRIN1) and 1 or more of the 4 NMDAR2 subunits: NMDAR2A (GRIN2A), NMDAR2B (GRIN2B), NMDAR2C (GRIN2C) and NMDAR2D (GRIN2D). Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Aug 2008]			

### - VALIDATION IMAGES -



Sample: Lane 1: Recombinant human GRIN2A protein, N-Trx-His(bs-42271P) Primary: Anti-GRIN2A (bs-3507R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 163 kDa Observed band size: 38 kDa

#### - SELECTED CITATIONS -

- [IF=13.352] Tingting Ku. et al. Tebuconazole mediates cognitive impairment via the microbe-gut-brain axis (MGBA) in mice. ENVIRON INT. 2023 Feb::107821 WB ;MOUSE. 36827814
- [IF=8.58] Ku, Tingting, et al. "NF-κB-regulated microRNA-574-5p underlies synaptic and cognitive impairment in response to atmospheric PM 2.5 aspiration." Particle and Fibre Toxicology 14.1 (2017): 34. Other ;="Mouse". 28851397
- **[IF=5.62]** Zhao, Tianyun, et al. "Ketamine administered to pregnant rats in the second trimester causes long-lasting behavioral disorders in offspring." Neurobiology of Disease (2014). WB ;="Rat". 24780497
- [IF=5.23] Ji, Ya-Bin, et al. "TFP5 peptide, derived from CDK5-activating cofactor p35, provides neuroprotection in earlystage of adult ischemic stroke." Scientific Reports 7 (2017): 40013. WB ;="MOUSe". 28045138

• [IF=5.1] Liang Xiao-Shan. et al. IRAK-M Ablation Promotes Status Epilepticus-Induced Neuroinflammation via Activating M1 Microglia and Impairing Excitatory Synaptic Function. MOL NEUROBIOL. 2023 Jun;:1-15 WB ;MOUSE. 37277682