
MMP13 Rabbit pAb

Catalog Number: bs-0575R

Target Protein: MMP13

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Human, Mouse, Rat (predicted:Rabbit)

Predicted MW: 52 kDa

Entrez Gene: 4322

Swiss Prot: P45452

Source: KLH conjugated synthetic peptide derived from human MMP13: 201-300/471.

Purification: affinity purified by Protein A


Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.


Background: bs-0575P is one synthetic peptide derived from human MMP13.

Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. The protein encoded by this gene cleaves type II collagen more efficiently than types I and III. It may be involved in articular cartilage turnover and cartilage pathophysiology associated with osteoarthritis. The gene is part of a cluster of MMP genes which localize to chromosome 11q22.3. [provided by RefSeq, Jul 2008].


VALIDATION IMAGES




Sample: Ovary (Rat) Lysate at 40 ug Primary: Anti- MMP13 (bs-0575R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 52 kD Observed band size: 52 kD




MG63 (Human) Lysate at 30 ug U2OS (Human) Lysate at 30 ug MDA-MB-231 (Human) Lysate at 30 ug Primary: Anti- MMP13 (bs-0575R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 52 kD Observed band size: 60 kD




Sample: Embryo (Mouse) Lysate at 40 ug Primary: Anti- MMP13 (bs-0575R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 52 kD Observed band size: 52 kD



Tissue/cell: rat articular cartilage; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-MMP-13 Polyclonal Antibody, Unconjugated (bs-0575R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody (SP-0023) and DAB (C-0010) staining



Tissue/cell: human bladder carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-MMP-13 Polyclonal Antibody, Unconjugated (bs-0575R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody (SP-0023) and DAB (C-0010) staining



Generously provided by Markus Linder from Medical University Vienna as part of the Bioss Discovery Program. Formalin-fixed, paraffin embedded, and decalcified in EDTA mouse bone labeled with Anti-MMP-13 Polyclonal Antibody, Unconjugated (bs-0575R) at 1:200 followed by conjugation to the secondary antibody and DAB staining

PRODUCT SPECIFIC PUBLICATIONS

[IF=10.684] Chen Zhang. et al. The novel hyaluronic acid granular hydrogel attenuates osteoarthritis progression by inhibiting the TLR-2/NF- κ B signaling pathway through suppressing cellular senescence. BIOENG TRANSL MED. 2022 Dec;;e10475 IHC ; Mouse . 10.1002/btm2.10475

[IF=8.8] Jiajun Chen. et al. Integrating UHPLC-MS/MS quantitative analysis and exogenous purine supplementation to elucidate the antidepressant mechanism of Chaigui granules by regulating purine metabolism. J PHARM ANAL. 2023 Aug;; WB ; Rat . 10.1016/j.jpha.2023.08.008

[IF=8.718] Yang, Jiashu. et al. m6A-mediated upregulation of AC008 promotes osteoarthritis progression through the miR-328-3p-AQP1/ANKH axis. Exp Mol Med. 2021 Nov;;1-12 WB ; Human . 34737423

[IF=6.832] Lingan Huang. et al. Zinc finger protein 521 attenuates osteoarthritis via the histone deacetylases 4 in the nucleus. BIOENGINEERED. 2022;13(6):14489-14502 IHC ; Rat . 36694467

[IF=6.832] Hou, Yonghui. et al. Nonwoven-based gelatin/polycaprolactone membrane loaded with ERK inhibitor U0126 for treatment of tendon defects. Stem Cell Res Ther. 2022 Dec;13(1):1-11 IHC ; Rat . 35012661