

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DERL3 Rabbit pAb

Catalog Number: bs-14281R

Target Protein: DERL3
Concentration: 1mg/ml

Form: Liquid Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500), ICC/IF (1:100-500), ELISA (1:5000-10000)

Reactivity: (predicted:Human, Mouse, Rat, Rabbit, Pig, Cow, Chicken)

Predicted MW: 27 kDa

Subcellular Cytoplasm

Locations:

Entrez Gene: 91319 Swiss Prot: Q96Q80

Source: KLH conjugated synthetic peptide derived from human DERL3: 51-150/235.

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: The protein encoded by this gene belongs to the derlin family, and resides in the

endoplasmic reticulum (ER). Proteins that are unfolded or misfolded in the ER must be refolded or degraded to maintain the homeostasis of the ER. This protein appears to be involved in the degradation of misfolded glycoproteins in the ER. Several alternatively spliced transcript variants encoding different isoforms have been identified for this gene.

[provided by RefSeq, Oct 2008]

PRODUCT SPECIFIC PUBLICATIONS

[IF=4.493] Qin Wentao. et al. Systematic Construction and Validation of a Novel Ferroptosis-Related Gene Model for Predicting Prognosis in Cervical Cancer. J IMMUNOL RES. 2022;2022:2148215 IHC; Human. 35935576

[IF=3.368] Manman Geng. et al. Up-regulated DERL3 in fibroblast-like synoviocytes exacerbates inflammation of rheumatoid arthritis. Clin Immunol. 2020 Nov;220:108579 IHC; Rat . 32866644