

bs-0295G-BF594

[Secondary Antibodies]

Goat Anti-Rabbit IgG H&L, BF594 conjugated

BioSS
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

| | | |
|---|---------------------|--------------------------------------|
| Host: Goat | Isotype: IgG | Applications: IF (1:200-1000) |
| Clonality: Polyclonal | | Flow-Cyt (1:50-200) |
| Target: Goat Anti-Rabbit IgG H&L | | ICC/IF (1:100-1000) |
| Purification: affinity purified by Protein G, nonspecific adsorbed | | Excitation Spectrum: 594nm |
| Concentration: 2.0 mg/ml | | Emission spectrum: 621nm |
| Storage: 10 mM TBS (pH=7.4) with 1% BSA, 0.03% Proclin300 and 50% glycerol. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. | | Reactivity: Rabbit |

— SELECTED CITATIONS —

- **[IF=14.7]** Xing Chen. et al. Genome-wide CRISPR screening identifies LRP1 as an entry factor for SFTSV. NAT COMMUN. 2025 Apr;16(1):1-13 IF ;Mouse. 40301361
- **[IF=8.2]** Lingling Yi. et al. NOD2 Promotes Sepsis-Induced Neuroinflammation by Increasing Brain Endoplasmic Reticulum Stress Mediated by LACC1. FREE RADICAL BIO MED. 2025 May;; IF ;Mouse. 40335000
- **[IF=4.6]** Feng Qu. et al. Synthesis and Biological Evaluation of Seco-Coumarin/Furoxan Hybrids as Potent Anti-Tumor Agents to Overcome Multidrug Resistance via Multiple Mechanisms. MOLECULES. 2025 May;30(11):2341 IF ;Human. 40509228
- **[IF=3.5]** MA Rui. et al. The glucose metabolism reprogramming of yak Sertoli cells under hypoxia is regulated by autophagy. BMC GENOMICS. 2025 Dec;26(1):1-18 IF ;Yak. 40251498
- **[IF=4.4]** Yan Yu. et al.hERG activators exhibit antitumor effects in breast cancer through calcineurin and β -catenin-mediated signaling pathways.FRONTIERS IN PHARMACOLOGY.2025 Jan 23;16:1545300. IF ;Rabbit. 39917621