

bs-0295G-BF594

[Secondary Antibodies]

Goat Anti-Rabbit IgG H&L, BF594 conjugated

Bioss
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Goat Clonality: Polyclonal Target: Goat Anti-Rabbit IgG H&L Purification: affinity purified by Protein G, nonspecific adsorbed Concentration: 2.0 mg/ml Storage: 10 mM TBS (pH=7.4) with 1% BSA, 0.03% Proclin300 and 50% glycerol. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.	Isotype: IgG	Applications: IF (1:200-1000) Flow-Cyt (1:50-200) Reactivity: Rabbit
---	---------------------	---

— SELECTED CITATIONS —

- **[IF=14.7]** Xing Chen. et al. Genome-wide CRISPR screening identifies LRP1 as an entry factor for SFTSV. NAT COMMUN. 2025 Apr;16(1):1-13 IF ;Mouse. 40301361
- **[IF=3.5]** Jing Wang. et al. The role of H3K27 acetylation in oxygen-glucose deprivation-induced spinal cord injury and potential for neuroprotective therapies. BRAIN RES BULL. 2025 Jan;220:111152 IF ;Rat. 39643249
- **[IF=3.5]** MA Rui. et al. The glucose metabolism reprogramming of yak Sertoli cells under hypoxia is regulated by autophagy. BMC GENOMICS. 2025 Dec;26(1):1-18 IF ;Yak. 40251498
- **[IF=4.4]** Yan Yu. et al.hERG activators exhibit antitumor effects in breast cancer through calcineurin and β -catenin-mediated signaling pathways.FRONTIERS IN PHARMACOLOGY.2025 Jan 23;16:1545300. IF ;Rabbit. 39917621
- **[IF=3.7]** Peng Liu. et al.Lactoferrin exhibits PEDV antiviral activity by interfering with spike-heparan sulfate proteoglycans binding and activating mucosal immune response.VETERINARY RESEARCH.2025 Jan 31;56(1):25. IF ;Goat. 39891300