

bs-0295G-BF594

[Secondary Antibodies]

Goat Anti-Rabbit IgG H&L, BF594 conjugated

BioSS
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Goat	Isotype: IgG	Applications: IF (1:200-1000)
Clonality: Polyclonal		Flow-Cyt (1:50-200)
Target: Goat Anti-Rabbit IgG H&L		ICC/IF (1:100-1000)
Purification: affinity purified by Protein G, nonspecific adsorbed		Excitation Spectrum: 594nm
Concentration: 2.0 mg/ml		Emission spectrum: 621nm
Storage: 10 mM TBS (pH=7.4) with 1% BSA, 0.03% Proclin300 and 50% glycerol. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Reactivity: Rabbit

— SELECTED CITATIONS —

- **[IF=14.7]** Xing Chen. et al. Genome-wide CRISPR screening identifies LRP1 as an entry factor for SFTSV. NAT COMMUN. 2025 Apr;16(1):1-13 IF ;Mouse. 40301361
- **[IF=13.2]** Yuan Su. et al. Oral solid-liquid biphasic microcapsules for precision targeted therapy of Salmonella-induced enteritis. CHEM ENG J. 2025 Apr;509:161497 IF ;Rat. 10.1016/j.cej.2025.161497
- **[IF=8.2]** Lingling Yi. et al. NOD2 Promotes Sepsis-Induced Neuroinflammation by Increasing Brain Endoplasmic Reticulum Stress Mediated by LACC1. FREE RADICAL BIO MED. 2025 May;; IF ;Mouse. 40335000
- **[IF=6.7]** Shi Liuliu. et al. Adeno-associated virus-mediated inhibition of ROCK2 promotes synaptogenesis and neurogenesis in rats after ischemic stroke. NEURAL REGEN RES. 2025 Jun;;10.4103/NRR.NRR IF ;Rat. 40537020
- **[IF=6.9]** Zhao Cui. et al. Identifying the target, mechanism, and agonist of α -ketoglutaric acid in delaying mesenchymal stem cell senescence. CELL REP. 2025 Jul;44: IF ;Human. 40570373