

bs-4865R**[Primary Antibody]****BNP Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

| | |
|---|---|
| Host: Rabbit Clonality: Polyclonal GeneID: 4879 Target: BNP Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: Brain natriuretic peptide (BNP), also known as B-type natriuretic peptide, is a hormone secreted by cardiomyocytes in the heart ventricles in response to stretching caused by increased ventricular blood volume. The 32-amino acid polypeptide BNP is secreted attached to a 76-amino acid N-terminal fragment in the prohormone called NT-proBNP (BNPT), which is biologically inactive. Once released, BNP binds to and activates the atrial natriuretic factor receptor NPRA, and to a lesser extent NPRB, in a fashion similar to atrial natriuretic peptide (ANP) but with 10-fold lower affinity. The biological half-life of BNP, however, is twice as long as that of ANP, and that of NT-proBNP is even longer, making these peptides better targets than ANP for diagnostic blood testing. | Isotype: IgG SWISS: P16860 Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Rat (predicted: Human, Mouse) Predicted MW.: 3/12 kDa Subcellular Location: Secreted |
|---|---|

— VALIDATION IMAGES —

Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-BNP Polyclonal Antibody, Unconjugated(bs-4865R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-BNP Polyclonal Antibody, Unconjugated(bs-4865R) 1:200, overnight at 4°C; The secondary antibody was Goat Anti-Rabbit IgG, Cy3 conjugated(bs-0295G-Cy3, red) used at 1:200 dilution for 40 minutes at 37°C. DAPI(5ug/ml, blue, C-0033) was used to stain the cell nuclei

— SELECTED CITATIONS —

- **[IF=4.069]** Zhao Q et al. CTRP15 derived from cardiac myocytes attenuates TGFβ1-induced fibrotic response in cardiac fibroblasts. Cardiovasc Drugs Ther. 2020 Oct;34(5):591-604. WB ;Mouse. 32424654
- **[IF=4.432]** Ruomiao Li. et al. Protective effects of dioscin against isoproterenol-induced cardiac hypertrophy via adjusting PKCε/ERK-mediated oxidative stress. Eur J Pharmacol. 2021 Sep;907:174277 WB ;Rat. 34171391
- **[IF=3.235]** Cao Z et al. AIN-93 Diet as an Alternative Model to Lieber-DeCarli Diet for Alcoholic Cardiomyopathy. Alcohol

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Clin Exp Res. 2019 Apr 29. IHC ;Mouse. 31034614

- **[IF=2.989]** Lin, Yueh-Min. et al. ZAK β Alleviates Oxidized Low-density Lipoprotein (ox-LDL)-Induced Apoptosis and B-type Natriuretic Peptide (BNP) Upregulation in Cardiomyoblast. CELL BIOCHEM BIOPHYS. 2022 Jul;;1-8 WB ;Rat. 35776316