bs-11440R

[Primary Antibody]

Bioss

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

THRB1 Rabbit pAb

- DATASHEET -

Host: Rabbit **Isotype:** IgG

Clonality: Polyclonal

GenelD: 7068 **SWISS:** P10828

Target: THRB1

Immunogen: KLH conjugated synthetic peptide derived from human Thyroid

Hormone Receptor beta: 201-300/461.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Thyroid hormone receptors (TRs) are ligand-dependent

transcription factors that mediate the biological activities of thyroid hormone (T3). Thyroid hormone receptor b2 (TRb2) is a high affinity receptor for triiodothyronine which belongs to the nuclear hormone receptor family and the NR1 subfamily. It is composed of three domains: a modulating N-terminal domain, a DNA-binding domain and a C-terminal steroid-binding domain. Defects in the receptor result in generalized thyroid hormone resistance (GTHR). GTHR is transmitted as an autosomal dominant trait, but an autosomal recessive form also exists. The disease is characterized by goiter, abnormal mental functions, increased susceptibility to infections, abnormal growth and bone maturation, tachycardia and deafness. GTHR patients also have high levels of circulating thyroid hormones (T3-T4), with normal or slightly elevated thyroid stimulating hormone.

Applications: WB (1:500-2000)

IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Flow-Cyt (1ug/Test)

Reactivity: Human, Mouse, Rat


(predicted: Rabbit, Sheep,

Cow, Chicken)


Predicted MW.: 53 kDa

Subcellular Location: Nucleus


VALIDATION IMAGES




Sample: Lane 1: Cerebrum (Mouse) Lysate at 40 ug Lane 2: Pituitary (Mouse) Lysate at 40 ug Lane 3: Adrenal gland (Mouse) Lysate at 40 ug Lane 4: Muscle (Mouse) Lysate at 40 ug Lane 4: Muscle (Mouse) Lysate at 40 ug Lane 6: Pituitary (Rat) Lysate at 40 ug Lane 6: Pituitary (Rat) Lysate at 40 ug Lane 7: Adrenal gland (Rat) Lysate at 40 ug Lane 8: Muscle (Rat) Lysate at 40 ug Lane 9: K562 (Human) Cell Lysate at 30 ug Lane 10: Siha (Human) Cell Lysate at 30 ug Lane 11: A431 (Human) Cell Lysate at 30 ug Lane 12: HepG2 (Human) Cell Lysate at 30 ug Primary: Anti-THRB1 (bs-11440R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 53/46 kD Observed band size: 53/46 kD




Paraformaldehyde-fixed, paraffin embedded (mouse liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (THRB1) Polyclonal Antibody, Unconjugated (bs-11440R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (THRB1) Polyclonal Antibody, Unconjugated (bs-11440R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Blank control: HepG2. Primary Antibody (green line): Rabbit Anti-THRB1 antibody (bs-11440R) Dilution: $1\mu g$ /10^6 cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody: Goat anti-rabbit IgG-PE Dilution: $1\mu g$ /test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at-20°C. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at at room temperature . Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.



Blank control: A431. Primary Antibody (green line): Rabbit Anti-THRB1 antibody (bs- 11440R) Dilution: 1ug/Test; Secondary Antibody: Goat anti-rabbit IgG-FITC Dilution: 0.5ug/Test.
Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

- SELECTED CITATIONS -

- [IF=4.8] Ying Zhang. et al. Long-term iodine deficiency and excess inhibit β-casein and α-lactalbumin secretion of milk in lactating rats. J NUTR BIOCHEM. 2024 Nov;:109812 WB ;Rat. 39603394
- [IF=4.8] Xiaoqing Shen. et al. Comparative effects of raw and processed cistanche glycosides on the HPT axis and gut microbiota in a rat model of kidney-yang deficiency. FRONT PHARMACOL. 2025 Jul;16: ;. 40786034
- [IF=4.9] Ying Zhang. et al. Negative Modulation of Maternal Iodine Deficiency and Excess on Milk Lipid Synthesis and Secretion in Lactating Rats. J NUTR BIOCHEM. 2025 Aug;:110085 WB;Rat. 40876516