[Primary Antibody]

phospho-Tau (Ser673) Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

– DATASHEET –		400-901-9800
Host: Rabbit Clonality: Polyclonal	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500)
GenelD: 4137	SWISS: P10636	IF (1:100-500)
Target: phospho-Tau (Ser673)		Reactivity: Human, Mouse, Rat (predicted: Rabbit, Cow, Chicken, Dog, Horse)
Immunogen: KLH conjugated Synthesised phosphopeptide derived from human Tau around the phosphorylation site of Ser673: IG(p-S)LD.		
Purification: affinity purified by Protein A		_
Concentration: 1mg/ml		Predicted MW.: ^{52/79} kDa
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Location: Cell membrane ,Cytoplasm
Background: Tau proteins are important Promotes microtubule assembly and stability, and might be involved in the establishment and maintenance of neuronal polarity. The C-terminus binds axonal microtubules while the N-terminus binds neural plasma membrane components, suggesting that tau functions as a linker protein between both. Axonal polarity is predetermined by tau localization (in the neuronal cell) in the domain of the cell body defined by the centrosome. The short isoforms allow plasticity of the cytoskeleton whereas the longer isoforms may preferentially play a role in its stabilization. Tau proteins subcellular located in the axons of neurons, in the cytoso I and in association with plasma membrane components. It expressed in neurons. PNS-tau is expressed in the central nervous system.		
- VALIDATION IMAGES -		

Ge^{40²}₀²⁰ ge⁴² 140 − 140 − 150 − 60 − − 35 − 35 −

Sample: Cerebrum (Rat) Lysate at 40 ug Liver (Mouse) Lysate at 40 ug HepG2(Human) Cell Lysate at 30 ug Primary: Anti- phospho-Tau (Ser673) (bs-10114R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 52/79 kD Observed band size: 60 kD