bs-8673R

[Primary Antibody]

KCTD12 Rabbit pAb

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 115207 SWISS: Q96CX2

Target: KCTD12

Immunogen: KLH conjugated synthetic peptide derived from human KCTD12:

251-325/325.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: The BTB (Broad-Complex, Tramtrack and Bric a brac) domain, also known as the POZ (POxvirus and Zinc finger) domain, is an Nterminal homodimerization domain that contains multiple copies of kelch repeats and/or C2H2-type zinc fingers. Proteins that contain BTB domains are thought to be involved in transcriptional regulation via control of chromatin structure and function. KCTD12 (potassium channel tetramerisation domain containing 12), also known as PFET1 or PFETIN, is a 325 amino acid protein that is expressed in fetal organs, with highest levels in the cochlea and brain and extremely low levels in adult organs, such as brain and lung. KCTD12 is considered a prognostic biomarker of

gastrointestinal stromal tumors.

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) **IF** (1:100-500) ICC/IF (1:100-500) **ELISA** (1:5000-10000)

Reactivity: (predicted: Human, Mouse,

Rat, Rabbit, Pig, Sheep, Cow, Dog, Horse)

Predicted MW.: 36 kDa

Subcellular Location: Cell membrane