

bs-1440R**[Primary Antibody]****Cytokeratin 5 Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

<p>Host: Rabbit</p> <p>Clonality: Polyclonal</p> <p>GeneID: 3852</p> <p>Target: Cytokeratin 5</p> <p>Immunogen: KLH conjugated synthetic peptide derived from human Cytokeratin 5: 451-580/580.</p> <p>Purification: affinity purified by Protein A</p> <p>Concentration: 1mg/ml</p> <p>Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.</p> <p>Background: Cytokeratins (CK) are intermediate filaments of epithelial cells, both in keratinising tissue (ie., skin) and non keratinising cells (ie., mesothelial cells). Although not a traditional marker for endothelial cells, cytokeratins have also been found in some microvascular endothelial cells. At least 20 different cytokeratins (CK) in the molecular range of 40 to 70 kDa and isoelectric points of 5 to 8.5 can be identified using two dimensional gel electrophoresis. Biochemically, most members of the CK family fall into one of two classes, type I (acidic polypeptides) and type II (basic polypeptides). At least one member of the acidic family and one member of the basic family is expressed in all epithelial cells. Defects in KRT5 are a cause of epidermolysis bullosa simplex.</p>	<p>Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)</p> <p>Reactivity: Human, Rat (predicted: Mouse, Rabbit, Pig, Cow, Dog, Horse)</p> <p>Predicted MW.: 64 kDa</p> <p>Subcellular Location: Extracellular matrix, Cytoplasm, Nucleus</p>
--	---

— VALIDATION IMAGES —

Sample: A431(Human) Cell Lysate at 40 ug
 Primary: Anti-Cytokeratin 5 (bs-1440R) at 1/300 dilution
 Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
 Predicted band size: 64kD
 Observed band size: 64 kD



Sample: Skin (Rat) Lysate at 40 ug
 Primary: Anti-Cytokeratin 5 (bs-1440R) at 1/1000 dilution
 Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
 Predicted band size: 64 kD
 Observed band size: 64kD