

---

## MRF/C11orf9 Rabbit pAb

Catalog Number: bs-11191R

Target Protein: MRF/C11orf9

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Human, Mouse, Rat (predicted:Rabbit, Pig, Sheep, Cow, Chicken, Dog, Horse)

Predicted MW: 124 kDa

Entrez Gene: 745

Source: KLH conjugated synthetic peptide derived from human MRF: 401-500/1151.

Purification: affinity purified by Protein A


Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.


**Background:** C11orf9 (chromosome 11 open reading frame 9), also known as MRF (myelin gene regulatory factor), Ndt80, KIAA0954 or MGC10781, is a 1,151 amino acid single-pass membrane protein that localizes to nucleus. A member of the MRF family, C11orf9 is expressed in the ARPE-19 cell line, brainstem, uterus and, to a lesser extent, in basal ganglion and liver. C11orf9 is weakly expressed in cerebellum and retina. C11orf9 contains one NDT80 DNA-binding domain and acts as a transcription factor that is required for expression of central nervous system (CNS) myelin genes such as MBP and MOG, thereby playing a central role in oligodendrocyte maturation and CNS myelination. C11orf9 exists as two alternatively spliced isoforms and is encoded by a gene located on human chromosome 11, which consists of approximately 135 million base pairs and 1,400 genes. Chromosome 11 makes up around 4% of human genomic DNA and is considered a gene and disease association dense chromosome.

### VALIDATION IMAGES


---




Sample: Lung (Mouse) Lysate at 40 ug Primary: Anti- MRF/C11orf9 (bs-11191R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 124 kD Observed band size: 124 kD



Sample: Lane 1: Human A549 cell lysates Lane 2: Human HepG2 cell lysates Lane 3: Human U2OS cell lysates Lane 3: Human A431 cell lysates Primary: Anti- MRF/C11orf9 (bs-11191R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 124 kDa Observed band size: 124 kDa



Sample: Uterus (Mouse) Lysate at 40 ug Primary: Anti- MRF/C11orf9 (bs-11191R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 124 kD Observed band size: 124 kD



Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-MRF/C11orf9 Polyclonal Antibody, Unconjugated(bs-11191R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

## PRODUCT SPECIFIC PUBLICATIONS

[IF=3.47] Xiaowei Yu. et al. Nanophthalmos-Associated MYRF Gene Mutation Causes Ciliary Zonule Defects in Mice. Invest Ophth Vis Sci. 2021 Mar;62(3):1-1 WB,IHC ; Mouse . 33646289