bs-1407R

[Primary Antibody]

Bioss

HIF1 beta Rabbit pAb

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 405 **SWISS:** P27540

Target: HIF1 beta

Immunogen: KLH conjugated synthetic peptide derived from human HIF-1 Beta:

681-789/789.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: This metastasis suppressor gene product is a membrane

glycoprotein that is a member of the transmembrane 4 superfamily. Expression of this gene has been shown to be downregulated in tumor progression of human cancers and can be activated by p53 through a consensus binding sequence in the promoter. Its expression and that of p53 are strongly correlated, and the loss of expression of these two proteins is associated with poor survival for prostate cancer patients. Two alternatively spliced transcript variants encoding distinct isoforms have been

found for this gene.

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) IF (1:100-500) Flow-Cyt (2ug/Test) ICC/IF (1:100-500)

Reactivity: Human, Mouse


(predicted: Rat, Rabbit, Pig,

Cow, Dog, Horse)


Predicted MW.: 87 kDa

Subcellular Location: Nucleus


VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (Mouse uterus); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HIF1 beta) Polyclonal Antibody, Unconjugated (bs-1407R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



HepG2 cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (HIF1 beta) polyclonal Antibody, Unconjugated (bs-1407R) 1:25, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.



Blank control (black line) :HepG2, Primary Antibody (green line): Rabbit Anti-HIF1 beta antibody (bs-1407R) Dilution: 2ug/Test: Secondary Antibody (white blue line): Goat anti-rabbit IgG-FITC Dilution: 0.5ug/Test. Isotype control (orange line) : Normal Rabbit IgG Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

- SELECTED CITATIONS -

• [IF=17.521] Huan Lei. et al. A Combination Therapy Using Electrical Stimulation and Adaptive, Conductive Hydrogels Loaded with Self-Assembled Nanogels Incorporating Short Interfering RNA Promotes the Repair of Diabetic Chronic

Wounds. Advanced Science. 2022 Sep;:2201425 IF; Rat. 36064844 • [IF=10.7] Taishan Liu. et al. Algae-inspired chitosan-pullulan-based multifunctional hydrogel for enhanced wound healing. CARBOHYD POLYM. 2025 Jan;347:122751 IF; Rat. 10.1016/j.carbpol.2024.122751