


**bsm-52401R****[ Primary Antibody ]****BioSS**  
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**MKLP1 Recombinant Rabbit mAb****DATASHEET****Host:** Rabbit**Clonality:** Recombinant**GeneID:** 9493**Target:** MKLP1**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.  
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.**Background:** The protein encoded by this gene is a member of kinesin-like protein family. This family includes microtubule-dependent molecular motors that transport organelles within cells and move chromosomes during cell division. This protein has been shown to cross-bridge antiparallel microtubules and drive microtubule movement in vitro. Alternate splicing of this gene results in two transcript variants encoding two different isoforms.**Isotype:** IgG**CloneNo.:** 7C9**SWISS:** Q02241**Applications:** WB (1:500-1000)**Flow-Cyt** (2ug/Test)**ICC/IF** (1:100-500)**Reactivity:** Human, Rat  
(predicted: Mouse)**Predicted  
MW.:** 110 kDa**Subcellular  
Location:** Cytoplasm ,Nucleus**VALIDATION IMAGES**

Sample: Lane 1: Rat Lymph node tissue lysates  
 Lane 2: Rat Cerebellum tissue lysates Lane 3: Rat  
 Testis tissue lysates Primary: Anti-MKLP1  
 (bsm-52401R) at 1/1000 dilution Secondary:  
 IRDye800CW Goat Anti-Rabbit IgG at 1/20000  
 dilution Predicted band size: 110 kDa Observed  
 band size: 100 kDa



Blank control:A549. Primary Antibody (green  
 line): Rabbit Anti-MKLP1 antibody (bsm-52401R)  
 Dilution: 2ul/Test; Secondary Antibody : Goat  
 anti-rabbit IgG-FITC Dilution: 0.5ug/Test.  
 Protocol The cells were fixed with 4% PFA  
 (10min at room temperature)and then  
 permeabilized with 90% ice-cold methanol for  
 20 min at room temperature.The cells were then  
 incubated in 5%BSA to block non-specific  
 protein-protein interactions for 30 min at room  
 temperature .Cells stained with Primary  
 Antibody for 30 min at room temperature. The  
 secondary antibody used for 40 min at room  
 temperature. Acquisition of 20,000 events was  
 performed.

**SELECTED CITATIONS**

- **[IF=13.417]** Kari L. Price. et al. Evolutionarily conserved midbody remodeling precedes ring canal formation during gametogenesis. DEV CELL. 2023 Mar 09 ICC ;Mouse. 36898376