


bsm-52401R**[Primary Antibody]****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

MKLP1 Recombinant Rabbit mAb**DATASHEET****Host:** Rabbit**Clonality:** Recombinant**GeneID:** 9493**Target:** MKLP1**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.**Background:** The protein encoded by this gene is a member of kinesin-like protein family. This family includes microtubule-dependent molecular motors that transport organelles within cells and move chromosomes during cell division. This protein has been shown to cross-bridge antiparallel microtubules and drive microtubule movement in vitro. Alternate splicing of this gene results in two transcript variants encoding two different isoforms.**Isotype:** IgG**CloneNo.:** 7C9**SWISS:** Q02241**Applications:** WB (1:500-1000)**Flow-Cyt** (2ug/Test)**ICC/IF** (1:50)**Reactivity:** Human, Rat
(predicted: Mouse)**Predicted
MW.:** 110 kDa**Subcellular
Location:** Cytoplasm ,Nucleus**VALIDATION IMAGES**


Sample: Lane 1: Rat Lymph node tissue lysates
Lane 2: Rat Cerebrum tissue lysates Lane 3: Rat Testis tissue lysates
Primary: Anti-MKLP1 (bsm-52401R) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 110 kDa
Observed band size: 100 kDa



A549 cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum) at 37°C for 20 min; Antibody incubation with (MKLP1) Monoclonal Antibody, Unconjugated (bsm-52401R) 1:50, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue) was used to stain the cell nuclei.



Hela cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum) at 37°C for 20 min; Antibody incubation with (MKLP1) Monoclonal Antibody, Unconjugated (bsm-52401R) 1:50, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue) was used to stain the cell nuclei.



Blank control: A549. Primary Antibody (green line): Rabbit Anti-MKLP1 antibody (bsm-52401R)
Dilution: 2ul/Test; Secondary Antibody: Goat anti-rabbit IgG-FITC Dilution: 0.5ug/Test.
Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at room temperature. The cells were then incubated in 5% BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

— SELECTED CITATIONS —

- **[IF=13.417]** Kari L. Price. et al. Evolutionarily conserved midbody remodeling precedes ring canal formation during gametogenesis. DEV CELL. 2023 Mar 09 ICC ;Mouse. 36898376