

**bsm-52305R****[ Primary Antibody ]****NFKB p65 Recombinant Rabbit mAb****Bioss**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**DATASHEET****Host:** Rabbit**Isotype:** IgG**Clonality:** Recombinant**CloneNo.:** 3G1**GeneID:** 5970**SWISS:** Q04206**Target:** NFKB p65**Immunogen:** A synthesized peptide derived from human NF-kB p65: 500-551/551.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

**Background:** NF-kappa-B is a ubiquitous transcription factor involved in several biological processes. It is held in the cytoplasm in an inactive state by specific inhibitors. Upon degradation of the inhibitor, NF-kappa-B moves to the nucleus and activates transcription of specific genes. NF-kappa-B is composed of NFKB1 or NFKB2 bound to either REL, RELA, or RELB. The most abundant form of NF-kappa-B is NFKB1 complexed with the product of this gene, RELA. Four transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2011].**Applications:** **WB** (1:500-5000)  
**IHC-P** (1:50-200)  
**IHC-F** (1:50-200)  
**IF** (1:50-200)  
**Flow-Cyt** (1:20-50)  
**ICC/IF** (1:50-200)**Reactivity:** Human, Mouse, Rat**Predicted MW.:** 61 kDa**Subcellular Location:** Cytoplasm ,Nucleus**VALIDATION IMAGES**


25 ug total protein per lane of various lysates (see on figure) probed with NFKB p65 monoclonal antibody, unconjugated (bsm-52305R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.




25 ug total protein per lane of various lysates (see on figure) probed with NFKB p65 monoclonal antibody, unconjugated (bsm-52305R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.




Paraformaldehyde-fixed, paraffin embedded Human Tonsil; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with NFKB p65 Monoclonal Antibody, Unconjugated(bsm-52305R) at 1:200 overnight at 4°C, followed by conjugation to the SP Kit (Rabbit, SP-0023) and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Colon Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with NFKB p65 Monoclonal Antibody, Unconjugated(bsm-52305R) at 1:200 overnight at 4°C, followed by conjugation to the SP Kit (Rabbit, SP-0023) and DAB (C-0010) staining.



4% Paraformaldehyde-fixed HeLa (left) cell and NFKB p65 knockout HeLa(right) cell; Triton X-100 at r.t. for 20 min; Antibody incubation with (NFKB p65) monoclonal Antibody, unconjugated (bsm-52305R) 1:100, 90 min at 37°C; followed by conjugated Goat Anti-Rabbit IgG antibody (green, bs-60295G-FITC) at 37°C for 90 min, DAPI (blue, C02-04002) was used to stain the cell




4% Paraformaldehyde-fixed HeLa (H) cell; Triton X-100 at r.t. for 20 min; Antibody incubation with (NFKB p65) monoclonal Antibody, unconjugated (bsm-52305R) 1:100, 90 min at 37°C; followed by conjugated Goat Anti-Rabbit IgG antibody (green, bs-60295G-FITC) at 37°C for 90 min, DAPI (blue, C02-04002) was used to stain the cell nuclei. PBS instead of the primary antibody was

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

nuclei.

used as the blank control.



The HeLa (H) cells were fixed with 4% PFA (10 min at r.t.) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, the cells then were incubated in 5% BSA to block non-specific protein-protein interactions (30 min at r.t.). Primary Antibody (green): Rabbit Anti-NFKB p65 antibody (bsm-52305R): 1 µg/10<sup>6</sup> cells; Secondary Antibody (white blue): Goat anti-Rabbit IgG-FITC (bs-60295G-FITC): 1 µg/test. Isotype Control (orange): Rabbit IgG (bs-0295P). Blank control (black): PBS. Acquisition of 20,000 events was performed.

## — SELECTED CITATIONS —

- **[IF=17.1]** Lei Liu. et al. Myricetin Oligomer Triggers Multi-Receptor Mediated Penetration and Autophagic Restoration of Blood-Brain Barrier for Ischemic Stroke Treatment. ACS NANO. 2024;XXXX(XXX):XXX-XXX WB ;Mouse. 38533773
- **[IF=5.6]** Yan-Qing Xie. et al. Anti-inflammatory and analgesic effects of Streblus indicus. FRONT PHARMACOL. 2023; 14: 1249234 WB ;Mouse. 37829300
- **[IF=5.6]** Yaxi Zhou. et al. Silkworm pupa protein peptide improved DSS-induced colitis in C57BL/6 mice through the MAPK/NF-κB signaling pathway. J FUNCT FOODS. 2023 Nov;110:105852 WB ;Mouse. 10.1016/j.jff.2023.105852
- **[IF=4.996]** Zhao, Lei. et al. Lactobacillus plantarum S9 alleviates lipid profile, insulin resistance, and inflammation in high-fat diet-induced metabolic syndrome rats. SCI REP-UK. 2022 Sep;12(1):1-10 WB ;Rat. 36109620
- **[IF=3.67]** Zhao Z et al. Lactobacillus plantarum NA136 ameliorates nonalcoholic fatty liver disease by modulating gut microbiota, improving intestinal barrier integrity, and attenuating inflammation. Appl Microbiol Biotechnol. 2020 Apr 26. WB ;Mouse. 32335723