bs-0403R-FITC

## [ Secondary Antibodies ]

## Rabbit Anti-MG IgG H&L, FITC conjugated



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

– DATASHEET –––––		400-901-9800
Host: Rabbit	<b>lsotype:</b> IgG	Applications: IF (1:100-1000)
Clonality: Polyclonal		Flow-Cyt (1:100-1000) ICC/IF (1:100-1000)
<b>Target:</b> Rabbit Anti-MG lgG H&L <b>Purification:</b> affinity purified by Protein A		Excitation Spectrum: 495nm Emission spectrum: 519nm
Concentration: 2.0 mg/ml		Reactivity: Mongolian gerbil
<b>Storage:</b> 10 mM TBS (pH=7.4) with 1% BSA, 0.03% Proclin300 and 50% glycerol. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our adaptive immune response. The IgG molecule has two separate functions; to bind to the pathogen that elicited the response and to recruit other cells and molecules to destroy the antigen. The variability of the IgG pool is generated by somatic recombination and the number of specificities in an individual at a given time point is estimated to be 1011 variants.		IgG e nd to on