

bs-0358D-FITC

[Secondary Antibodies]



www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

Donkey Anti-Guinea Pig IgG H&L, FITC conjugated

— DATASHEET —

<p>Host: Donkey</p> <p>Clonality: Polyclonal</p> <p>Target: Donkey Anti-Guinea Pig IgG H&L</p> <p>Purification: affinity purified by Protein G</p> <p>Concentration: 2.0 mg/ml</p> <p>Storage: 10 mM TBS (pH=7.4) with 1% BSA, 0.03% Proclin300 and 50% glycerol. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.</p> <p>Background: Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our adaptive immune response. The IgG molecule has two separate functions; to bind to the pathogen that elicited the response and to recruit other cells and molecules to destroy the antigen. The variability of the IgG pool is generated by somatic recombination and the number of specificities in an individual at a given time point is estimated to be 1011 variants.</p>	<p>Isotype: IgG</p> <p>Applications: IF (1:100-1000) Flow-Cyt (1:100-1000) ICC/IF (1:100-1000)</p> <p>Excitation Spectrum: 495nm Emission spectrum: 519nm</p> <p>Reactivity: Guinea Pig</p>
--	---

— SELECTED CITATIONS —

- **[IF=4.8]** Hai-bo Liu. et al. Dispelling Dampness, Relieving Turbidity and Dredging Collaterals Decoction, Attenuates Potassium Oxonate-Induced Hyperuricemia in Rat Models. DRUG DES DEV THER. 2023 Aug 01 WB ;Rat. 37551408