bs-6471R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

ITPR3 Rabbit pAb

GenelD: 3710 SWISS: Q14573

Target: ITPR3

Immunogen: KLH conjugated synthetic peptide derived from human ITPR3:

21-120/2671.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Inositol 1,4,5-triphosphate (IP3) functions as a second messenger for a myriad of extracellular stimuli including hormones, growth factors and neurotransmitters. Receptor tyrosine kinases indirectly increase the intracellular levels of IP3 through the activation of phospholipases such as phospholipase C (PLC), which convert phosphatidylinositol-4,5 bisphosphate into IP3 and diacylglycerol (DAG). The inositol 1,4,5-triphosphate receptor, IP3R, acts as an inositol triphosphate (IP3)-gated calcium release channel in a variety of cell types. Three IP3 receptor subtypes have been described and are designated IP3R-I, IP3R-II and IP3R-III. IP3R-I is the predominant IP3R subtype expressed in neuronal tissues and the central nervous system, but is also expressed at high levels in the liver.

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) **IF** (1:50-200)

Reactivity: Mouse (predicted: Human,


Rat, Rabbit, Cow, Chicken,

Dog, Horse)

Predicted 294 kDa

Subcellular Location: Cell membrane ,Cytoplasm

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (mouse cerebellum); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ITPR3) Polyclonal Antibody, Unconjugated (bs-6471R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

- SELECTED CITATIONS -

- [IF=5.8] Lin Yiqun. et al. Mitochondria-associated endoplasmic reticulum membrane as a mediator of vanadiuminduced endoplasmic reticulum quality control in duck brains. ENVIRON SCI POLLUT R. 2024 Mar;:1-17 IF; Duck. 38446297
- [IF=4.155] Junjun Peng. et al. Endoplasmic reticulum-mitochondria coupling attenuates vanadium-induced apoptosis

laying and laying periods using an iTRAQ-based approach." PLOS ONE 12.9 (2017): e0185253. WB ;Other Species. 28945779						