

bs-9880R**[Primary Antibody]****ELYS Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

Host: Rabbit Clonality: Polyclonal GeneID: 25909 Target: ELYS Immunogen: KLH conjugated synthetic peptide derived from human ELYS: 451-550/2266. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: ELYS is a protein of the nuclear matrix that contains one AT hook DNA-binding domain. ELYS is a dual nucleoporin/kinetochore protein required for nuclear pore assembly and proper cell division. Nuclear pores span the nuclear envelope and act as gated aqueous channels to regulate the transport of macromolecules between the nucleus and cytoplasm, from individual proteins and RNAs to entire viral genomes. The A.T. hook near the C-terminus of likely indicates ELYS can function as a transcription factor which may play a specific role in hematopoietic tissues. It also may play a role in early embryo development, as ELYS is critical for the survival of inner cells of the blastocyst.	Isotype: IgG SWISS: Q8IZA4	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Mouse (predicted: Human, Rat, Pig, Sheep, Cow, Horse) Predicted MW.: 252 kDa Subcellular Location: Cytoplasm ,Nucleus
---	---	--

— VALIDATION IMAGES —

Sample: Spinal cord (Mouse) Lysate at 40 ug
Primary: Anti- ELYS (bs-9880R) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 252 kD
Observed band size: 252 kD



Paraformaldehyde-fixed, paraffin embedded (mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ELYS) Polyclonal Antibody, Unconjugated (bs-9880R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.