

**bs-9612R****[ Primary Antibody ]****Bioss**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**PIBF1 Rabbit pAb****— DATASHEET —**


<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> IHC-P (1:100-500) <b>IHC-F</b> (1:100-500) <b>IF</b> (1:100-500) <b>ICC/IF</b> (1:100)  <b>Reactivity:</b> Human, Rat (predicted: Mouse, Rabbit, Sheep, Cow, Dog)  <b>Predicted MW.:</b> 90 kDa  <b>Subcellular Location:</b> Secreted ,Nucleus
<b>Clonality:</b> Polyclonal		
<b>GeneID:</b> 10464	<b>SWISS:</b> Q8WXW3	
<b>Target:</b> PIBF1		
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from human PIBF1/C13orf24: 551-651/757.		
<b>Purification:</b> affinity purified by Protein A		
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> C13orf24 is also known as Progesterone-induced-blocking factor 1 (PIBF1). PIBF1 is synthesized by lymphocytes of healthy pregnant women in the presence of progesterone and inhibits arachidonic acid release, by acting on the phospholipase A2 enzyme, inducing a Th2 biased immune response. It also controls NK activity, and modifies the cytokine balance exerting an anti-abortion effect.		

**— VALIDATION IMAGES —**

Paraformaldehyde-fixed, paraffin embedded (rat spleen); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (PIBF1) Polyclonal Antibody, Unconjugated (bs-9612R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human breast carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PIBF1) Polyclonal Antibody, Unconjugated (bs-9612R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Hela cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (PIBF1) polyclonal Antibody, Unconjugated (bs-9612R) 1:100, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.