


**bs-20778R****[ Primary Antibody ]****Angiotensinogen Rabbit pAb****BioSS**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**— DATASHEET —****Host:** Rabbit**Isotype:** IgG**Clonality:** Polyclonal**Target:** Angiotensinogen**Immunogen:** KLH conjugated synthetic peptide derived from mouse Angiotensinogen: 101-200/477.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.  
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.**Background:** bs-0587P is a Eight branched multiple antigenic peptide of Angiotensin II.**Applications:** **WB** (1:500-2000)**IHC-P** (1:100-500)**IHC-F** (1:100-500)**IF** (1:100-500)**Reactivity:** Mouse, Rat  
(predicted: Human)**Predicted**  
**MW.:** 49 kDa**Subcellular**  
**Location:** Secreted**— VALIDATION IMAGES —**

Sample: Lane 1: Plasma (Mouse) at 20 ug Lane 2: Kidney (Mouse) Lysate at 40 ug Lane 3: Adrenal gland (Mouse) Lysate at 40 ug Lane 4: Spleen (Rat) Lysate at 40 ug Lane 5: Plasma (Rat) at 20 ug Primary: Anti-Angiotensinogen (bs-20778R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 50-70 kD Observed band size: 60 kD



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Angiotensinogen) Polyclonal Antibody, Unconjugated (bs-20778R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.