

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

phospho-Tau (Ser198) Rabbit pAb

Catalog Number: bs-4199R

Target Protein: phospho-Tau (Ser198)

Concentration: 1mg/ml

Form: Liquid Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: Flow-Cyt (1µg/Test)

Reactivity: Human, Mouse, Rat (predicted:Rabbit, Dog, Horse)

Predicted MW: 52/79 kDa

Entrez Gene: 4137 Swiss Prot: P10636

Source: KLH conjugated Synthesised phosphopeptide derived from human Tau around the

phosphorylation site of Ser198: GY(P-S)SP.

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Tau proteins are important Promotes microtubule assembly and stability, and might be

involved in the establishment and maintenance of neuronal polarity. The C-terminus binds axonal microtubules while the N-terminus binds neural plasma membrane components,

suggesting that tau functions as a linker protein between both. Axonal polarity is


predetermined by tau localization (in the neuronal cell) in the domain of the cell body defined by the centrosome. The short isoforms allow plasticity of the cytoskeleton whereas

the longer isoforms may preferentially play a role in its stabilization. Tau proteins

subcellular located in the axons of neurons, in the cytoso I and in association with plasma membrane components. It expressed in neurons. PNS-tau is expressed in the peripheral

nervous system while the others are expressed in the central nervous system.

VALIDATION IMAGES



Blank control (blue line): MCF 7 (fixed with 70% methanol (Overnight at 4°C) and then permeabilized with 90% ice-cold methanol for 30 min on ice). Primary Antibody (green line): Rabbit Anti- phospho-Tau protein (Ser198)antibody (bs-4199R),Dilution: $0.2\mu g / 10^5$ cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody (white blue line): Goat anti-rabbit IgG-FITC,Dilution: $1\mu g / test$.