bs-0335R-FITC

## [ Secondary Antibodies ]

## Rabbit Anti-Monkey IgG H&L, FITC conjugated



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

| - DATASHEET  |   | 400-901-9800  |
|--|---|---|
| Host: Rabbit<br>Clonality: Polyclonal  | <b>Isotype:</b> IgG   | Applications: IF (1:100-1000)<br>Flow-Cyt (1:100-1000)  |
| Target: Rabbit Anti-Monkey IgG H&L Purification: affinity purified by Protein A  |   | ICC/IF (1:100-1000)<br>Excitation Spectrum: 495nm<br>Emission spectrum: 519nm<br>Reactivity: Monkey |
| glycerol.  | with 1% BSA, 0.03% Proclin300 and 50<br>year. Avoid repeated freeze/thaw cyc  | 1%  |
| Background: Immunoglobulin G (I<br>serum with normal le<br>is important for our c<br>molecules are produ<br>adaptive immune res<br>functions; to bind to<br>recruit other cells an<br>variability of the IgG | gG), is one of the most abundant prote<br>wels between 8-17 mg/mL in adult blo<br>lefence against microorganisms and th<br>ced by B lymphocytes as a part of our<br>ponse. The IgG molecule has two sepa<br>the pathogen that elicited the response<br>d molecules to destroy the antigen. Th<br>pool is generated by somatic recombin<br>pecificities in an individual at a given ti | eins in<br>bod. IgG<br>he<br>arate<br>se and to<br>ne<br>nation                                     |