

bsm-33114M**[Primary Antibody]****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

Histone H3(di methyl K9) Mouse mAb**— DATASHEET —**

Host: Mouse	Isotype: IgG2b	Applications: WB (1:500-2000) ELISA (1:1000-5000) Reactivity: Human, Mouse, Rat (predicted: Cow) Predicted MW.: 15 kDa Subcellular Location: Nucleus
Clonality: Monoclonal	CloneNo.: 5D2	
GeneID: 8350	SWISS: P68431	
Target: Histone H3(di methyl K9)		
Immunogen: KLH conjugated synthesised methylpeptide derived from human Histone H3 around the methylation site of di methyl K9: AR(Di Methyl-K)ST.		
Purification: affinity purified by Protein G		
Concentration: 1mg/ml		
Storage: Size : 50ul/100ul/200ul 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Size : 200ug (PBS only) 0.01M PBS Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Modulation of the chromatin structure plays an important role in the regulation of transcription in eukaryotes. The nucleosome, made up of four core histone proteins (H2A, H2B, H3 and H4), is the primary building block of chromatin. The N-terminal tail of core histones undergoes different posttranslational modifications including acetylation, phosphorylation and methylation. These modifications occur in response to cell signal stimuli and have a direct effect on gene expression. In most species, the histone H2B is primarily acetylated at lysines 5, 12, 15 and 20. Histone H3 is primarily acetylated at lysines 9, 14, 18 and 23. Acetylation at lysine 9 appears to have a dominant role in histone deposition and chromatin assembly in some organisms. Phosphorylation at Ser10 of histone H3 is tightly correlated with chromosome condensation during both mitosis and meiosis.		

— VALIDATION IMAGES —

25 ug total protein per lane of various lysates
(see on figure) probed with Histone H3(di methyl
K9) monoclonal antibody, unconjugated
(bsm-33114M) at 1:1000 dilution and 4°C
overnight incubation. Followed by conjugated
secondary antibody incubation at r.t. for 60 min.