## bs-11838R

## [ Primary Antibody ]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

## Emx1 Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**GenelD: 2016 SWISS:** Q04741

Target: Emx1

**Immunogen:** KLH conjugated synthetic peptide derived from human Emx1:

201-257/257.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.


Background: Emx1 and Emx2 are human homologs to the Drosophila

developmental genes empty spiracles expressed in anterior body regions during early Drosophila embryogenesis. Emx1 and Emx2 are homeobox proteins expressed in the developing vertebrate brain. Emx2 is expressed in the dorsal telencephalon and small diencephalic regions, while Emx1 expression is exclusively confined to pyramidal neurons of the dorsal telencephalon. In the embryonic brain, Emx1 is expressed in both proliferating and differentiating neurons while Emx2 is expressed only in proliferating neurons. OTX1 and OTX2 are human homologs of the Drosophila developmental genes orthodenticle. In development, the sequence of expression begins with OTX2 at day 10 post coitum followed by OTX1, Emx2 and finally Emx1. The genes encoding human Emx1 and Emx2 map to chromosomes 2p14-p13 and 10q26.1, respectively.

VALIDATION IMAGES



Sample: HepG2 Cell (Human) Lysate at 30 ug Primary: Anti- Emx1 (bs-11838R)at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 29kD Observed band size: 29kD



Blank control: HepG2. Primary Antibody (green line): Rabbit Anti-Emx1 antibody (bs-11838R) Dilution: 1µg/10^6 cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody: Goat anti-rabbit IgG-PE Dilution: 1µg /test, Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at-20°C. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

Flow-Cyt (1ug/Test)

Reactivity: Human (predicted: Mouse,

Rat, Rabbit, Pig, Sheep, Cow, Chicken, Dog, Horse)

Predicted 29 kDa

MW.:

Subcellular Nucleus