bs-11841R

[Primary Antibody]

BIOSS ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

FNBP1L Rabbit pAb

GeneID: 54874 SWISS: Q5T0N5

Target: FNBP1L

Immunogen: KLH conjugated synthetic peptide derived from human

FNBP1L/TOCA1: 101-200/605.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: The protein encoded by this gene binds to both CDC42 and N-

WASP. This protein promotes CDC42-induced actin polymerization by activating the N-WASP-WIP complex and, therefore, is involved

in a pathway that links cell surface signals to the actin

cytoskeleton. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul

2008]

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) IF (1:100-500)


Reactivity: Human (predicted: Mouse,

Rat, Rabbit, Pig, Sheep, Cow, Chicken, Dog, Horse)

Predicted MW.: 70 kDa

Subcellular Location: Cell membrane ,Cytoplasm

VALIDATION IMAGES



Tissue/cell: human liver cancer; 4%
Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-FNBP1L Polyclonal Antibody, Unconjugated(bs-11841R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining