bs-11604R

- DATASHEET -

[Primary Antibody]

Isotype: IgG

Glyt1 Rabbit pAb

Host: Rabbit



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Applications: WB (1:500-2000)

Reactivity: Human, Mouse, Rat (predicted: Rabbit, Cow, Dog)

Predicted MW.: ^{78 kDa}

Subcellular Location: Cell membrane

Clonality: Polyclonal GenelD: 6536 Target: Glyt1 Immunogen: KLH conjugated synthetic peptide derived from human Glyt1/SLC6A9: 209-285/706. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: Na+/Cl- dependent neurotransmitter transporters are a superfamily of transmembrane proteins that contain 12 membrane spanning regions (1). Specifically, the highly hydrophobic Na+/Cldependent glycine transporters (GlyT) are crucial for the termination of neurotransmission at glycinergic synapses (2,3). Two different GlyT genes encode GlyT2 and GlyT1, which exists as two isoforms produced by alternative splicing of the same gene located on human chromosome 1p31.3 (3,4). The GlyT1 gene may be an early marker of neural development and encodes gliaspecific transporter proteins (3). Although GlyT1 and GlyT2 are both expressed in the brain and spinal cord, each shows a unique pattern of expression (3,5,6). GlyT1 is found only in the white matter of the CNS, whereas GlyT2 is found in the gray matter of the CNS as well as in macrophages and mast cells in peripheral tissues (3,5). The anatomic distribution of GlyT2 mRNA suggests that glycine may act as a supraspinal neurotransmitter and may function as a chemical messenger outside the CNS (5).

- VALIDATION IMAGES



Sample: Cerebrum (Mouse) Lysate at 40 ug Cerebrum(Rat) Lysate at 40 ug Primary: Anti-Glyt1 (bs-11604R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 78 kD Observed band size: 78 kD