

bs-14381R**[Primary Antibody]****DNAJC14 Rabbit pAb**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500) ELISA (1:5000-10000) Reactivity: (predicted: Human, Mouse, Rat, Horse) Predicted MW.: 79 kDa Subcellular Location: Cytoplasm
Clonality: Polyclonal		
GeneID: 85406	SWISS: Q6Y2X3	
Target: DNAJC14		
Immunogen: KLH conjugated synthetic peptide derived from human DNAJC14: 151-250/702.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: The DnaJ family is one of the largest of all the chaperone families and has evolved with diverse cellular localization and functions. The presence of the J domain defines a protein as a member of the DnaJ family. DnaJ heat shock induced proteins are from the bacterium Escherichia coli and are under the control of the htpR regulatory protein. The DnaJ proteins play a critical role in the HSP 70 chaperone machine by interacting with HSP 70 to stimulate ATP hydrolysis. DnaJ proteins are important mediators of proteolysis and are involved in the regulation of protein degradation, exocytosis and endocytosis. DnaJC14 (DnaJ homolog subfamily C member 14), also known as DRIP78 (Dopamine receptor-interacting protein of 78 kDa) and HDJ3 (Human DnaJ protein 3), is a 702 amino acid endoplasmic reticular membrane protein that contains one J domain. DnaJC14 regulates the export of target proteins, such as the dopamine D1 receptor (D1DR), from the endoplasmic reticulum to the cell surface.		