

bs-3353R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

phospho-PIM1 (Tyr309) Rabbit pAb**— DATASHEET —**

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500)
Clonality: Polyclonal		IHC-F (1:100-500)
GeneID: 5292	SWISS: P11309	IF (1:100-500)
Target: PIM1 (Tyr309)		Reactivity: Human, Mouse, Rat (predicted: Rabbit, Pig, Sheep, Cow, Dog)
Immunogen: KLH conjugated Synthesised phosphopeptide derived from human PIM1 around the phosphorylation site of Tyr309: HR(p-Y)HG.		Predicted MW.: 45 kDa
Purification: affinity purified by Protein A		Subcellular Location: Cell membrane ,Cytoplasm ,Nucleus
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: PIM1 is a member of the serine/threonine kinase PIM oncogene family. PIM1 has been implicated in lymphomagenesis, cell proliferation, apoptosis, differentiation and tumourigenesis. The PIM1 protein kinase is upregulated in prostate cancer. The Pim family serine/threonine protein kinases were first identified in studies examining genes targeted for proviral insertion in murine leukemia virus-induced T lymphomas. Increased levels of Pim kinases predispose cells to lymphomagenesis and enhance the activity of mitogenic proteins such as p100, c-Myb, and Cdc25A. In addition, Pim kinases are also involved in modulation of synaptic strength in neurons and anti-apoptotic signaling in hematopoietic progenitor cells. Pim-3, a member of the proto-oncogene Pim family that expresses serine/threonine kinase activity, shares significant homology with Pim-1 serine/threonine protein kinases. Pim-3 may function as a mediator of synaptic plasticity in the brain and is presumably involved in the anti-apoptosis process and cell cycle progression as well as the proliferation of human hepatoma cell lines. The Pim-3 protein is widely expressed, however no expression is observed in the colon, thymus, or small intestine.		

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (human rectal carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Phospho-PIM1 (Tyr309)) Polyclonal Antibody, Unconjugated (bs-3353R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Phospho-PIM1) Polyclonal Antibody, Unconjugated (bs-3353R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.