
human CD274 (B7-H1, PD-L1) Mouse mAb

Catalog Number: bsm-30139M

Target Protein: human CD274 (B7-H1, PD-L1)

Form: Size : 100ug

Liquid

Size : 200ug (PBS only)

Lyophilized

Note: Centrifuge tubes before opening. Reconstitute the lyophilized product in distilled water. Optimal concentration should be determined by the end user.

Host: Mouse

Clonality: Monoclonal

Clone No.: 4C11

Isotype: IgG1

Applications: Flow-Cyt (1ug/Test)

Reactivity: Human

Predicted MW: 32 kDa

Detected MW: 40 kDa

Subcellular: Secreted ,Extracellular matrix

Locations:

Purification: affinity purified by Protein G

Storage: Size : 100ug

0.01M PBS (pH7.4).

Size : 200ug (PBS only)


0.01M PBS

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: This gene encodes an immune inhibitory receptor ligand that is expressed by hematopoietic and non-hematopoietic cells, such as T cells and B cells and various types of tumor cells. The encoded protein is a type I transmembrane protein that has immunoglobulin V-like and C-like domains. Interaction of this ligand with its receptor inhibits T-cell activation and cytokine production. During infection or inflammation of normal tissue, this interaction is important for preventing autoimmunity by maintaining homeostasis of the immune response. In tumor microenvironments, this interaction provides an immune escape for tumor cells through cytotoxic T-cell inactivation. Expression of this gene in tumor cells is considered to be prognostic in many types of human malignancies, including colon cancer

and renal cell carcinoma. Alternative splicing results in multiple transcript variants.
[provided by RefSeq, Sep 2015]

VALIDATION IMAGES



Blank control (black line) :MDA-MB-231. Primary Antibody (green line): Mouse Anti-Human PD-L1 antibody (bsm-30139M) Dilution:1ug/Test; Secondary Antibody (white blue line) : Goat anti-mouse IgG-FITC Dilution: 0.5ug/Test. Isotype control (orange line) : Normal Mouse IgG Protocol The cells were incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.