

**bs-0604R****[ Primary Antibody ]****BioSS**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**Galectin 9 Rabbit pAb****— DATASHEET —**

<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> IHC-P (1:100-500)
<b>Clonality:</b> Polyclonal		<b>IHC-F</b> (1:100-500)
<b>GeneID:</b> 16859	<b>SWISS:</b> O08573	<b>IF</b> (1:100-500)
<b>Target:</b> Galectin 9		<b>Reactivity:</b> Human, Mouse, Rat
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from mouse galectin 9: 101-180/353.		
<b>Purification:</b> affinity purified by Protein A		<b>Predicted MW.:</b> 39 kDa
<b>Concentration:</b> 1mg/ml		<b>Subcellular Location:</b> Secreted ,Cytoplasm
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> Binds galactosides. May play a role in thymocyte-epithelial interactions relevant to the biology of the thymus. Is a ligand for HAVCR2/TIM3. Induces T-helper type 1 lymphocyte (Th1) death (By similarity). May provide the molecular basis for urate flux across cell membranes, allowing urate that is formed during purine metabolism to efflux from cells and serving as an electrogenic transporter that plays an important role in renal and gastrointestinal urate excretion. Highly selective to the anion urate.		

**— VALIDATION IMAGES —**

Paraformaldehyde-fixed, paraffin embedded (Human tonsil); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (galectin 9) Polyclonal Antibody, Unconjugated (bs-0604R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human colon carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (galectin 9) Polyclonal Antibody, Unconjugated (bs-0604R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

**— SELECTED CITATIONS —**

- **[IF=9.7]** Peier Chen. et al. Targeted heart repair by Tβ4-loaded cardiac-resident macrophage-derived extracellular vesicles modified with monocyte membranes. ACTA BIOMATER. 2023 Aug.; WB ;Mouse. 37597679
- **[IF=3.47]** Zhang Y et al. Konjac glucomannan improves hyperuricemia through regulating xanthine oxidase, adenosine deaminase and urate transporters in rats. Journal of Functional Foods,2018 48, 566–575. WB ;Rat. 10.1016/j.jff.2018.07.062
- **[IF=2.1]** Hui Zhao. et al. Expression of Tim-3/Galectin-9 pathway and CD8+T cells and related factors in patients with

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

cystic echinococcosis. EXP PARASITOL. 2023 Nov;254:108623 IHC ;Human. 37793539