

**bs-14394R****[ Primary Antibody ]****DNAJC5G Rabbit pAb****BioSS**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**— DATASHEET —**

<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)
<b>Clonality:</b> Polyclonal		
<b>GeneID:</b> 285126	<b>SWISS:</b> Q8N7S2	
<b>Target:</b> DNAJC5G		<b>Reactivity:</b> Mouse, Rat (predicted: Human, Rabbit)
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from human DNAJC5G: 101-189/189.		
<b>Purification:</b> affinity purified by Protein A		<b>Predicted MW.:</b> 21 kDa
<b>Concentration:</b> 1mg/ml		<b>Subcellular Location:</b> Cell membrane
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> The DnaJ family is one of the largest of all the chaperone families and has evolved with diverse cellular localization and functions. Members of the DnaJ family are characterized by the presence of the J domain. DnaJ heat shock induced proteins are from the bacterium Escherichia coli and are under the control of the htpR regulatory protein. The DnaJ proteins play a critical role in the HSP 70 chaperone machine by interacting with HSP 70 to stimulate ATP hydrolysis. Proteins of the DnaJ family contain cysteine rich regions that are composed of zinc fingers that form a peptide binding domain responsible for chaperone function. DnaJ proteins are important mediators of proteolysis and are involved in the regulation of protein degradation, exocytosis and endocytosis.		

**— VALIDATION IMAGES —**

Paraformaldehyde-fixed, paraffin embedded (Rat testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (DNAJC5G) Polyclonal Antibody, Unconjugated (bs-14394R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (DNAJC5G) Polyclonal Antibody, Unconjugated (bs-14394R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.