

**bsm-33103M**

**[ Primary Antibody ]**

## CK18 Mouse mAb

**Bioss**  
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

### — DATASHEET —

**Host:** Mouse

**Clonality:** Monoclonal

**GeneID:** 3875

**Target:** CK18

**Purification:** affinity purified by Protein G

**Concentration:** 1mg/ml

**Storage:** Size : 50ul/100ul/200ul  
0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.  
Size : 200ug (PBS only)  
0.01M PBS  
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

**Background:** KRT18 encodes the type I intermediate filament chain keratin 18. Keratin 18, together with its filament partner keratin 8, are perhaps the most commonly found members of the intermediate filament gene family. They are expressed in single layer epithelial tissues of the body. Mutations in this gene have been linked to cryptogenic cirrhosis. Two transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Jul 2008]

**Isotype:** IgG

**CloneNo.:** 10H2

**SWISS:** P05783


**Applications:** WB (1:1000-2000)  
IHC-P (1:200-1000)  
IHC-F (1:200-1000)  
IF (1:200-1000)  
Flow-Cyt (1ug/Test)  
ICC/IF (1:100-500)

**Reactivity:** Human


**Predicted MW.:** 48 kDa

**Subcellular Location:** Cytoplasm ,Nucleus


### — VALIDATION IMAGES —




Sample: Lane 1: HeLa (Human) cell Lysate at 40 ug  
Lane 2: HepG2 (Human) cell Lysate at 40 ug  
Lane 3: A549 (Human) cell Lysate at 40 ug  
Lane 4: MCF-7 (Human) cell Lysate at 40 ug  
Primary: Anti-CK18 (bsm-33103M) at 1/1000 dilution  
Secondary: IRDye800CW Goat Anti-Mouse IgG at 1/20000 dilution  
Predicted band size: 48 kD  
Observed band size: 45 kD




Paraformaldehyde-fixed, paraffin embedded (human pancreatic cancer); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CK18) Monoclonal Antibody, Unconjugated (bsm-33103M) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Mouse)(sp-0024) instructions and DAB staining.




Paraformaldehyde-fixed, paraffin embedded (human gastric carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CK18) Monoclonal Antibody, Unconjugated (bsm-33103M) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Mouse)(sp-0024) instructions and DAB staining.



Tissue/cell: HeLa cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum,



Paraformaldehyde-fixed, paraffin embedded (human pancreatic carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for



The HeLa (H) cells were fixed with 4% PFA (10 min at r.t.) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, the cells then

**Important Note:** This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

C-0005) at 37°C for 20 min; Antibody incubation with (CK18) monoclonal Antibody, Unconjugated (bsm-33103M) 1:100, 90 minutes at 37°C; followed by a CY3 conjugated Goat Anti-Mouse IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.

15min; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CK18) Monoclonal Antibody, Unconjugated (bsm-33103M ) at 1:500 overnight at 4°C, followed by a conjugated Goat Anti- Mouse IgG antibody (YF488) for 90 minutes, and DAPI for nuclei staining.

were incubated in 5%BSA to block non-specific protein-protein interactions (30 min at r.t.), followed by secondary antibody incubation for 40 min at room temperature. Primary Antibody (green):Mouse Anti-CK18 antibody (bsm-33103M): 1 µg/10<sup>6</sup> cells; Isotype Control (orange): Mouse IgG (bs-0296P). Blank control (black): PBS. Acquisition of 20,000 events was performed.

---

## — SELECTED CITATIONS —

- **[IF=4.9]** Bao Yuan. et al. Exploring the Mechanism of H<sub>2</sub>S Synthesis in Male Bactrian Camel Poll Glands Based on Data Independent Acquisition Proteomics and Non-Targeted Metabolomics. INT J MOL SCI. 2024 Jul; 25(14): 7700 IF ;Camel. 39062942
- **[IF=4.4]** Wang Bin. et al. Enhancing endometrial receptivity: the roles of human chorionic gonadotropin in autophagy and apoptosis regulation in endometrial stromal cells. REPROD BIOL ENDOCRIN. 2024 Dec;22(1):1-18 IF ;Human. 38576003