- DATASHEET -----

[Primary Antibody]

Cathelicidin/Camp Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASTILL			
Host:	Rabbit	lsotype: IgG	Applications: IHC-P (1:100-500)
Clonality:	Polyclonal	IHC-F (1:100-500) IF (1:100-500)	
GenelD:	12796	SWISS: P51437	ICC/IF (1:100-500)
Target:	Cathelicidin/Camp		ELISA (1:5000-10000)
Immunogen: KLH conjugated synthetic peptide derived from mouse Camp: 101-170/170.			Reactivity: (predicted: Mouse, Rat)
Purification:	affinity purified by Pro	tein A	
Concentration:	1mg/ml		Predicted
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.			Subcellular Location: Secreted
peroxidase-negative granules of neutrophils. Along with the family of proteins known as defensins, cathelicidins participate in the first line of defense by preventing local infection and systemic invasion of microbes. FALL-39 precursor (FALL-39 peptide antibiotic, cationic anti-microbial protein, CAMP, CAP-18, HSD26) is a cathelicidin anti-microbial protein that contains the antibacterial peptide LL-37 (amino acids 134-170). In contrast to the defensins, which are cysteine-rich peptide that fold in \int -pleated sheets, LL-37 is a cysteine-free peptide that can adopt an amphipathic å- helical conformation. LL-37 binds to bacterial lipopolysaccharides (LPS) and is a potent chemotactic factor for recruiting mast cells to sites of inflammation. LL-37 is present in inflammatory skin diseases that include psoriasis, sub-acute lupus erthematosus, dermatitis and nickel contact hypersensitivity. It is not found in normal skin epidermis. The secreted protein is expressed primarily in bone marrow, testis and neutrophils. The mouse and rat ortholog, CRAMP (cathelin-related antimicrobial peptide), is also part of the cathelicidin family of host defense peptides. These include precursors of potent antimicrobial peptides that direct antimicrobial activity against various microbial pathogens and also activate mesenchymal cells during wound repair. CRAMP is expressed in testis, spleen, stomach and intestine. This gene encodes a member of an antimicrobial peptide family, characterized by a highly conserved N- terminal signal peptide containing a cathelin domain and a structurally variable cationic antimicrobial peptide, which is produced by extracellular proteolysis from the C-terminus. The protein plays an important role in innate immunity defense against viruses. In addition to its antibacterial, antifungal, and antiviral activities, the encoded protein functions in cell chemotaxis, immune mediator induction, and		y sh so ly so q,	

- SELECTED CITATIONS -

- [IF=5.1] Xinyue Qi. et al. Probiotics suppress LL37 generated rosacea-like skin inflammation by modulating
- TLR2/MyD88/NF-κB signaling pathway. FOOD FUNCT. 2024 Jul;: IHC ;MOUSE. 39143863
- [IF=5.3] Jun Zhu. et al. Eicosatrienoic acid inhibits estradiol synthesis through the CD36/FOXO1/CYP19A1 signaling

pathway to improve PCOS in mice. BIOCHEM PHARMACOL. 2024 Sep;:116517 WB ;MOUSE. 39236935