

bs-14410R**[Primary Antibody]****DOPEY2 Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)
Clonality: Polyclonal		
GeneID: 9980	SWISS: Q9Y3R5	
Target: DOPEY2		Reactivity: Rat (predicted: Human, Mouse, Pig, Sheep, Cow, Dog, Horse)
Immunogen: KLH conjugated synthetic peptide derived from human DOPEY2: 51-150/2298.		
Purification: affinity purified by Protein A		Predicted MW.: 258 kDa
Concentration: 1mg/ml		Subcellular Location: Secreted ,Extracellular matrix ,Cytoplasm
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Dopey-2 is a 2,298 amino acid protein that is ubiquitously expressed with high levels found in the developing central nervous system where it is thought to play a role in protein trafficking between early endosomes and the late Golgi. Multiple isoforms of Dopey-2 exist due to alternative splicing events. The gene encoding Dopey-2 maps to human chromosome 21 and may be involved in the pathogenesis of Down syndrome. The smallest of the human chromosomes, chromosome 21 comprises about 1.5% of the human genome and contains nearly 300 genes and 47 million base pairs. Down syndrome, also known as trisomy 21, is the disease most commonly associated with chromosome 21. Alzheimer's disease, Jervell and Lange-Nielsen syndrome and amyotrophic lateral sclerosis are also associated with chromosome 21. Translocations are found to occur between chromosome 21 and 8, and chromosome 21 and 12, in certain leukemias.		

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (DOPEY2) Polyclonal Antibody, Unconjugated (bs-14410R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.