bs-13401R

[Primary Antibody]

Bioss ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

phospho-AMPK alpha 1 (Ser496) Rabbit pAb

- DATASHEET -

Host: Rabbit **Isotype:** IgG

Clonality: Polyclonal

GeneID: 105787 **SWISS:** Q5EG47

Target: AMPK alpha 1 (Ser496)

Immunogen: KLH conjugated Synthesised phosphopeptide derived from mouse

AMPK alpha 1 around the phosphorylation site of Ser496: SG(p-

S)IS.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: The protein encoded by this gene belongs to the ser/thr protein

kinase family. It is the catalytic subunit of the 5'-prime-AMP-activated protein kinase (AMPK). AMPK is a cellular energy sensor conserved in all eukaryotic cells. The kinase activity of AMPK is activated by the stimuli that increase the cellular AMP/ATP ratio. AMPK regulates the activities of a number of key metabolic enzymes through phosphorylation. It protects cells from stresses that cause ATP depletion by switching off ATP-consuming biosynthetic pathways. Alternatively spliced transcript variants encoding distinct isoforms have been observed. [provided by

RefSeq, Jul 2008]

Applications: WB (1:500-2000)


IHC-P (1:100-500) **IHC-F** (1:100-500) **IF** (1:100-500)

Reactivity: Rat (predicted: Mouse)


Predicted MW.: 64 kDa

Subcellular Location: Cytoplasm ,Nucleus

VALIDATION IMAGES



Sample: Skeltal muscl(Rat) Cell Lysate at 40 ug Primary: Anti-phospho-AMPK alpha 1 (Ser496) (bs-13401R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 64 kD Observed band size: 64 kD



Paraformaldehyde-fixed, paraffin embedded (Rat skeletal muscle); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-AMPK alpha 1 (Ser496)) Polyclonal Antibody, Unconjugated (bs-13401R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

— SELECTED CITATIONS —

• [IF=3.7] Sun Xuri. et al. Exogenous NT-3 Promotes Phenotype Switch of Resident Macrophages and Improves Sciatic Nerve Injury through AMPK/NF-kB Signaling Pathway. NEUROCHEM RES. 2024 Jun;:1-15 WB; Rat. 38904909