

bs-13402R**[Primary Antibody]****phospho-AMPK alpha 1 (Ser496) Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

Host: Rabbit Clonality: Polyclonal GeneID: 5562 Target: AMPK alpha 1 (Ser496) Immunogen: KLH conjugated Synthesised phosphopeptide derived from human AMPK alpha 1 around the phosphorylation site of Ser496: SG(p-S)VS. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: The protein encoded by this gene belongs to the ser/thr protein kinase family. It is the catalytic subunit of the 5'-prime-AMP-activated protein kinase (AMPK). AMPK is a cellular energy sensor conserved in all eukaryotic cells. The kinase activity of AMPK is activated by the stimuli that increase the cellular AMP/ATP ratio. AMPK regulates the activities of a number of key metabolic enzymes through phosphorylation. It protects cells from stresses that cause ATP depletion by switching off ATP-consuming biosynthetic pathways. Alternatively spliced transcript variants encoding distinct isoforms have been observed. [provided by RefSeq, Jul 2008]	Isotype: IgG Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Human, Mouse Predicted MW.: 64 kDa Subcellular Location: Cytoplasm ,Nucleus
---	---

— VALIDATION IMAGES —

Sample: MCF-7(Human) Cell Lysate at 40 ug
PANC-1(Human) Cell Lysate at 40 ug Primary:
Anti-phospho-AMPK alpha 1 (Ser496)
(bs-10196R) at 1/300 dilution Secondary:
IRDye800CW Goat Anti-Rabbit IgG at 1/20000
dilution Predicted band size: 64 kD Observed
band size: 64 kD



Paraformaldehyde-fixed, paraffin embedded
(Human brain glioma); Antigen retrieval by
boiling in sodium citrate buffer (pH6.0) for
15min; Block endogenous peroxidase by 3%
hydrogen peroxide for 20 minutes; Blocking
buffer (normal goat serum) at 37°C for 30min;
Antibody incubation with (phospho-AMPK alpha
1 (Ser496)) Polyclonal Antibody, Unconjugated
(bs-13402R) at 1:400 overnight at 4°C, followed
by operating according to SP Kit(Rabbit)
(sp-0023) instructions and DAB staining.

— SELECTED CITATIONS —

- **[IF=4.225]** Li H et al . Trimetazidine Ameliorates Myocardial Metabolic Remodeling in Isoproterenol-Induced Rats Through Regulating Ketone Body Metabolism via Activating AMPK and PPAR α Front Pharmacol. 2020 Aug 14;11:1255. Other ;. 32922293
- **[IF=3.063]** Mengmeng Yu. et al. Taurine attenuates gossypol-induced apoptosis of C2C12 mouse myoblasts via the

GPR87-AMPK/AKT signaling. Amino Acids. 2020 Sep;52(9):1285-1298 WB ;Mouse. 32918616

- **[IF=3.1]** Tian Xiaorong. et al. Heme Oxygenase-1-Modified BMSCs Activate AMPK–Nrf2–FTH1 to Reduce Severe Steatotic Liver Ischemia–Reperfusion Injury. DIGEST DIS SCI. 2023 Sep;:1-16 IF,WB ;Rat. 37707747