## bs-20771R

## [ Primary Antibody ]

## **TERT Rabbit pAb**



sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

– DATASHEF	т ———		400-901-9800
Host:	Rabbit	Isotype: IgG	Applications: WB (1:500-2000
Clonality:	Polyclonal		Flow-Cyt (1µg, ICC/IF (1:100)
GenelD:	7015	SWISS: 014746	
Target:	TERT		Reactivity: Human, Mouse (predicted: Rat
Immunogen:	KLH conjugated synthetic 521-620/1132.	c peptide derived from human TERT:	
Purification: affinity purified by Protein A			Predicted
Concentration:	1mg/ml		MW.:
Storage:	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Location: <sup>Cytoplasm</sup> ,Nue
<b>Background:</b> Telomerase is a ribonucleoprotein polymerase that maintains telomere ends by addition of the telomere repeat TTAGGG. The enzyme consists of a protein component with reverse transcriptase activity, encoded by this gene, and an RNA component which serves as a template for the telomere repeat. Telomerase expression plays a role in cellular senescence, as it is normally repressed in postnatal somatic cells resulting in progressive shortening of telomeres. Deregulation of telomerase expression in somatic cells may be involved in oncogenesis. Studies in mouse suggest that telomerase also participates in chromosomal repair, since de novo synthesis of telomere repeats may occur at double-stranded breaks. Alternatively spliced variants encoding different isoforms of telomerase reverse transcriptase have been identified; the full-length sequence of some variants has not been determined. Alternative splicing at this locus is thought to be one mechanism of regulation of telomerase activity. [provided by RefSeq, Jul 2008].			

## - VALIDATION IMAGES



Sample: RAW264.7(Mouse) Cell Lysate at 30 ug Primary: Anti-TERT (bs-20771R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 124 kD Observed band size: 135 kD



Sample: Jurkat(Human) Cell Lysate at 30 ug RAW264.7(Mouse) Cell Lysate at 30 ug Hela(Human) Cell Lysate at 30 ug Primary: Anti-TERT (bs-20771R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 124 kD Observed band size: 135 kD



Hela cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (TERT) polyclonal Antibody, Unconjugated (bs-20771R) 1:100, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.







0) /Test)

, Dog)

cleus

Tissue/cell:A549 cell; 4% Paraformaldehydefixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Antibody incubation with (TERT) polyclonal Antibody, Unconjugated (bs-20771R) 1:100, 90 minutes at 37°C; followed by a FITC conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei. Blank control (blue line): Mouse thymus cells (fixed with 70% methanol (Overnight at 4°C) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C). Primary Antibody (green line): Rabbit Anti-TERT antibody (bs-20771R),Dilution: 1µg /10^6 cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody (white blue line): Goat antirabbit IgG-PE,Dilution: 1µg /test. Blank control:Hela. Primary Antibody (green line): Rabbit Anti-TERT antibody (bs-20771R) Dilution: 1µg /10^6 cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody : Goat anti-rabbit IgG-AF647 Dilution: 1µg /test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at-20°C. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature . Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature.