
CD19 Rabbit pAb

Catalog Number: bs-20782R

Target Protein: CD19

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: Flow-Cyt (1µg/Test)

Reactivity: Human, Mouse (predicted:Rat, Rabbit, Pig, Sheep, Cow, Horse)

Predicted MW: 59 kDa

Entrez Gene: 930

Swiss Prot: P15391

Source: KLH conjugated synthetic peptide derived from human CD19: 431-500/556.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: This gene encodes a member of the immunoglobulin gene superfamily. Expression of this cell surface protein is restricted to B cell lymphocytes. This protein is a reliable marker for pre-B cells but its expression diminishes during terminal B cell differentiation in antibody secreting plasma cells. The protein has two N-terminal extracellular Ig-like domains separated by a non-Ig-like domain, a hydrophobic transmembrane domain, and a large C-terminal cytoplasmic domain. This protein forms a complex with several membrane proteins including complement receptor type 2 (CD21) and tetraspanin (CD81) and this complex reduces the threshold for antigen-initiated B cell activation. Activation of this B-cell antigen receptor complex activates the phosphatidylinositol 3-kinase signalling pathway and the subsequent release of intracellular stores of calcium ions. This protein is a target of chimeric antigen receptor (CAR) T-cells used in the treatment of lymphoblastic leukemia. Mutations in this gene are associated with the disease common variable immunodeficiency 3 (CVID3) which results in a failure of B-cell differentiation and impaired secretion of immunoglobulins. CVID3 is characterized by hypogammaglobulinemia, an inability to mount an antibody response to antigen, and recurrent bacterial infections. Alternative splicing results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq, Jul 2020]

VALIDATION IMAGES



Blank control (blue line): HL60 cells (fixed with 70% methanol (Overnight at 4°C)). Primary Antibody (green line): Rabbit Anti-CD19 antibody (bs-20782R), Dilution: 1µg /10⁶ cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody (white blue line): Goat anti-rabbit IgG-PE, Dilution: 1µg /test.

PRODUCT SPECIFIC PUBLICATIONS

[IF=2.515] Li T et al. Digital gene expression analyses of mammary glands from meat ewes naturally infected with clinical mastitis. R. Soc. open sci. 2019,6: 181604. IHC ; Sheep . doi:10.1098/rsos.181604