### bs-13671R

### [ Primary Antibody ]

# BIOSS ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

## TRAF6BP Rabbit pAb

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 8887 SWISS: Q86VP1

Target: TRAF6BP

Immunogen: KLH conjugated synthetic peptide derived from human TRAF6BP:

241-340/789.

**Purification:** affinity purified by Protein A

Concentration: 1mg/ml

**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Inhibits TNF-induced apoptosis by mediating the TNFAIP3 anti-

apoptotic activity. Degraded by caspase-3-like family proteins upon TNF-induced apoptosis. May also play a role in the pro-

inflammatory cytokine IL-1 signaling cascade.

**Applications: IHC-P** (1:100-500)

**IHC-F** (1:100-500) **IF** (1:100-500)

Reactivity: Mouse (predicted: Human,

Rat, Sheep, Cow, Dog,


Horse)

Predicted MW.: 91 kDa

MW.:

**Subcellular Location:** Cell membrane ,Cytoplasm

### VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (mouse lung); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (TRAF6BP) Polyclonal Antibody, Unconjugated (bs-13671R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

#### - SELECTED CITATIONS -

• [IF=7.169] Wang Xuan-zhong. et al. TAX1BP1 contributes to deoxypodophyllotoxin-induced glioma cell parthanatos via inducing nuclear translocation of AIF by activation of mitochondrial respiratory chain complex I. ACTA PHARMACOL SIN. 2023 Apr;:1-14 WB,ICC; Human. 37186123