

bs-9901R**[Primary Antibody]****PAFAHG Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500)
Clonality: Polyclonal		IHC-F (1:100-500)
GeneID: 5050	SWISS: Q15102	IF (1:100-500)
Target: PAFAHG		Reactivity: Mouse, Rat (predicted: Human, Rabbit, Pig, Sheep, Cow, Dog, Horse)
Immunogen: KLH conjugated synthetic peptide derived from human PAFAH1B3/PAFAHG: 1-100/231.		Predicted MW.: 26 kDa
Purification: affinity purified by Protein A		Subcellular Location: Cytoplasm
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: The platelet activating factor (PAF) acetylhydrolases catalyze hydrolysis of the sn-2 ester bond of PAF and related pro-inflammatory phospholipids and thus attenuate their bioactivity. The family of PAF acetylhydrolases includes one secreted plasma isozyme and two intracellular isozymes. The intracellular isozymes are distinguished by differences in their primary sequence, tissue localization, subunit composition and substrate preferences. The most thoroughly characterized intracellular isoform, PAFAH1B, is a heterotrimeric protein expressed in brain tissue and plays an important role in brain development and function. PAFAH1B is comprised of a regulatory subunit (LIS1) and two homologous (63% identity) catalytic subunits (PAFAH1B2 and PAFAH1B3), which harbor all the activity of the enzyme. The PAFAH1B2 and PAFAH1B3 subunits readily associate with very high affinity to form heterodimers, and this dimerization is essential for both stability and catalytic activity. PAFAH1B3 is also commonly known as PAFAH1B 29kDa subunit, PAFAH1B subunit or PAFAH1B subunit α 1.		

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PAFAHG) Polyclonal Antibody, Unconjugated (bs-9901R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PAFAHG) Polyclonal Antibody, Unconjugated (bs-9901R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

— SELECTED CITATIONS —

- **[IF=0]** Vandenberghe L T M et al. Platelet-activating factor acetylhydrolase 1B3 (PAFAH1B3) is required for the formation

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

of the meiotic spindle during invitro oocyte maturation.Reproduction, Fertility and Development.2018. WB,ICC
;Bovine, Murine&Human. 10.1071/rd18019