bs-9536R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Geminin Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GeneID: 51053 **SWISS:** 075496

Target: Geminin

Immunogen: KLH conjugated synthetic peptide derived from human Geminin:

111-209/209.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Inhibits DNA replication by preventing the incorporation of MCM

complex into prereplication complex (pre-RC). It is degraded during the mitotic phase of the cell cycle. Its destruction at the metaphase-anaphase transition permits replication in the

succeeding cell cycle.

Geminin is a nuclear protein that regulates the initiation of DNA replication during the cell cycle. DNA replication requires the coordinated association of cdc6 and minichromosome maintenance (MCM) proteins with chromatin. Geminin blocks this assembly of the MCM into the prereplication complex and, in turn, prevents replication from occurring. Expression of geminin fluctuates throughout the cell cycle with geminin levels lowest at G1. Throughout S, G2 and M phases, geminin levels are consistently elevated followed by a decrease during mitosis. The initiation of DNA replication is dependent on the the degradation of geminin during mitosis and the absence of geminin throughout G1 phase. Geminin degradation is mediated by the anaphasepromoting complex (APC), which specifically targets B-type cyclins and other proteins containing a destruction box motif for degradation by ubiquitin-mediated proteolysis.

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) **IF** (1:100-500) Flow-Cyt (1µg/Test)

Reactivity: Human, Mouse

(predicted: Rat, Rabbit, Pig, Cow, Chicken, Dog, Horse)


Predicted 23 kDa

Subcellular Location: Cytoplasm ,Nucleus

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (Mouse small intestine); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Geminin) Polyclonal Antibody, Unconjugated (bs-9536R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Blank control(blue): Hela cells(fixed with 2% paraformaldehyde (10 min), then permeabilized with 90% ice-cold methanol for 30 min on ice). Primary Antibody:Rabbit Anti-Geminin antibody(bs-9536R), Dilution: $1\mu g$ in $100~\mu L$ 1XPBS containing 0.5% BSA; Isotype Control Antibody: Rabbit IgG(orange) ,used under the same conditions); Secondary Antibody: Goat anti-rabbit IgG-PE(white blue), Dilution: 1:200 in 1 X PBS containing 0.5% BSA.