## bs-3699R

## [ Primary Antibody ]

# www.bioss.com.cn

sales@bioss.com.cn techsupport@bioss.com.cn

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**GPR15 Rabbit pAb** 

**GeneID: 2838 SWISS:** P49685

Target: GPR15

Immunogen: KLH conjugated synthetic peptide derived from human GPR15:

51-150/360. < Extracellular >

Purification: affinity purified by Protein A

Concentration: 1mg/ml

**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: This gene encodes a G protein-coupled receptor that acts as a

chemokine receptor for human immunodeficiency virus type 1 and 2. The encoded protein localizes to the cell membrane. [provided  $\,$ 

by RefSeq, Nov 2012]

Applications: IHC-P (1:100-500)

400-901-9800

IHC-F (1:100-500) **IF** (1:200-800)

Reactivity: Mouse, Rat

(predicted: Human, Rabbit,

Dog, Horse)


Predicted 40 kDa

**Subcellular Location:** Cell membrane ,Cytoplasm

### **VALIDATION IMAGES**



Tissue/cell: Mouse stomach tissue; 4% Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer ( 0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-GPR15 Polyclonal Antibody, Unconjugated(bs-3699R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: rat stomach tissue; Frozen section; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-GPR15 Polyclonal Antibody, Unconjugated(bs-3699R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining