bs-0358G-PE

[Secondary Antibodies]

Goat Anti-Guinea Pig IgG H&L, PE conjugated



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

– DATASHEET –		400-901-9800
Host: Goat	Isotype: IgG	Applications: IF (1:100-1000)
Clonality: Polyclonal		Flow-Cyt (1:100-1000) ICC/IF (1:100-1000) Excitation Spectrum: 496/564nm Emission spectrum: 578nm
Target: Goat Anti-Guinea Pig IgG H&L		
Purification: affinity purified by Protein G Concentration: 2.0 mg/ml		Reactivity: Guinea Pig
glycerol.	with 1% BSA, 0.03% Proclin300 and 50% e year. Avoid repeated freeze/thaw cycles.	
serum with normal le is important for our o molecules are produ adaptive immune res functions; to bind to recruit other cells an variability of the IgG	gG), is one of the most abundant proteins in evels between 8-17 mg/mL in adult blood. IgG lefence against microorganisms and the ced by B lymphocytes as a part of our sponse. The IgG molecule has two separate the pathogen that elicited the response and t d molecules to destroy the antigen. The pool is generated by somatic recombination becificities in an individual at a given time be 1011 variants.	

- SELECTED CITATIONS -

• [IF=5.714] Haiwei Cheng. et al. Promoting immunity with novel targeting antigen delivery vehicle based on bispecific nanobody. INT IMMUNOPHARMACOL. 2023 Jun;119:110140 FCM ;Mouse. 37116343