

**bs-11675R****[ Primary Antibody ]****UBXD2 Rabbit pAb****Bioss**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**— DATASHEET —**


<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> <b>WB</b> (1:500-2000) <b>IHC-P</b> (1:100-500) <b>IHC-F</b> (1:100-500) <b>IF</b> (1:100-500)  <b>Reactivity:</b> Human, Mouse, Rat (predicted: Pig, Sheep, Cow, Dog, Horse)  <b>Predicted MW.:</b> 57 kDa  <b>Subcellular Location:</b> Cell membrane ,Cytoplasm
<b>Clonality:</b> Polyclonal		
<b>GeneID:</b> 23190	<b>SWISS:</b> Q92575	
<b>Target:</b> UBXD2		
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from human UBXD2: 201-300/508.		
<b>Purification:</b> affinity purified by Protein A		
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> Erasin is an endoplasmic reticulum (ER) and nuclear envelope membrane protein. Expressed in a variety of tissues, such as brain, placenta, heart, liver, prostate, kidney, pancreas, lung and skeletal muscle, erasin contains one UBX domain and participates in the clearing of ERAD (endoplasmic reticulum-associated protein degradation) substrates. The UBX domain of erasin is responsible for mediating its direct interaction with VCP (valosin-containing protein), an AAA-ATPase molecular chaperone. In response to ER stress, erasin expression is induced. The knockdown of erasin expression leads to the inhibition of ERAD, suggesting an important function of erasin in the ERAD pathway. In addition, erasin may be involved in Alzheimer' s disease, as it is known to accumulate in neurofibrillary degenerating neurons in patients with Alzheimer' s disease.		

**— VALIDATION IMAGES —**

Sample: Heart (Mouse) Lysate at 40 ug Cerebrum (Rat) Lysate at 40 ug  
Primary: Anti-UBXD2 (bs-11675R) at 1/1000 dilution  
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution  
Predicted band size: 57 kD Observed band size: 57 kD



Paraformaldehyde-fixed, paraffin embedded (Human kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (UBXD2) Polyclonal Antibody, Unconjugated (bs-11675R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (UBXD2) Polyclonal Antibody, Unconjugated (bs-11675R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.